

Review

Natural antifouling compound production by microbes associated with marine macroorganisms – A review



Sathianeson Satheesh*, Mohammad Abdulaziz Ba-akdah, Abdulmohsin A. Al-Sofyani

Department of Marine Biology, Faculty of Marine Sciences, King Abdulaziz University, Jeddah, Saudi Arabia

ARTICLE INFO

Article history:

Received 25 October 2015

Accepted 20 January 2016

Available online 18 February 2016

Keywords:

Antifouling

Biofouling

Bioactive compounds

Marine biotechnology

Marine microbes

ABSTRACT

In the marine environment, all hard surfaces including marine macroorganisms are colonized by microorganisms mainly from the surrounding environment. The microorganisms associated with marine macroorganisms offer tremendous potential for exploitation of bioactive metabolites. Biofouling is a continuous problem in marine sectors which needs huge economy for control and cleaning processes. Biotechnological way for searching natural product antifouling compounds gained momentum in recent years because of the environmental pollution associated with the use of toxic chemicals to control biofouling. While, natural product based antifoulants from marine organisms particularly sponges and corals attained significance due to their activities in field assays, collection of larger amount of organisms from the sea is not a viable one. The microorganisms associated with sponges, corals, ascidians, seaweeds and seagrasses showed strong antimicrobial and also antifouling activities. This review highlights the advances in natural product antifoulants research from microbes associated with marine organisms.

© 2016 Pontificia Universidad Católica de Valparaíso. Production and hosting by Elsevier B.V. All rights reserved.

Contents

1. Introduction	26
2. Eco-friendly antifoulants from marine organisms	27
3. Marine macroorganisms–microbe associations	28
4. Antifouling activities of microbes associated with marine invertebrates and ascidians	28
5. Antifouling activities of microbes associated with seaweeds and seagrasses 5	29
6. <i>Pseudoalteromonas</i> associated with marine organisms: A potential genus for antifouling research	30
7. Advantages of microbes as a source of bioactive metabolites	30
8. Conclusions and future perspectives	31
Conflict of interest statement	32
References	32

1. Introduction

Biofouling (accumulation of organisms) is a common problem on man-made objects submerged in the marine waters throughout the world. The biofouling growth on a substratum in the aquatic environment is a complex process (Fig. 1) with initial biofilm formation (consisting of microbes and microalgae) followed by settlement of

invertebrate larvae and algal spores [1,2]. Biofouling assemblage in marine environment is made up of thousands of marine organisms such as bacteria, fungi, phytoplankton, polychaetes, barnacles, molluscs, ascidians and algae (Fig. 2). Biofouling on submerged surfaces in the marine environment has considerable ecological and economical importance particularly serious implications for shipping, offshore aquaculture and coastal industries [3,4,5]. The effects are mainly due to the loss of productivity in aquaculture [6] or increased costs of fuel to shipping as well as the costs associated with ongoing prevention, management and control [7,8].

Due to the economic significance of the problem in the marine waters, various control strategies are adopted by the marine sectors

* Corresponding author.

E-mail addresses: ssathianeson@kau.edu.sa, satheesh_s2005@yahoo.co.in

(S. Satheesh).

Peer review under responsibility of Pontificia Universidad Católica de Valparaíso.




Fig. 1. Progression of biofouling development on hard substratum submerged in tropical coastal waters.

(Fig. 3). Tributyltin (TBT) containing antifouling paints were widely used in the commercial vessels to control biofouling [9,10]. However, use of TBT caused environmental problems as it is more toxic to non-target marine organisms [11,12,13,14]. Due to the environmental concern over the use of TBT, the International Maritime Organization and Marine Environment Protection Committee banned the application of TBT for marine applications from January 1, 2008 [5]. After the ban of TBT containing antifoulants, copper paints are used as an alternative despite the higher toxicity of copper to the marine environment [15]. In addition, many other compounds are also commonly used as antifouling biocides [16,17] which include irgarol 1051 diuron, dichloroflumid, chlorothalonil, zine, pyriothione, pyridine and zineb [10]. Some of these antifouling compounds are now under strict regulations in various regions due to the possible effects on marine ecosystems. Natural products are suggested as an alternative to toxic biocides in the antifouling paints for controlling biofouling. In a previous review by Qian et al. [10] have highlighted the recent progress in natural product antifouling research which consists of both

marine and terrestrial sources. Recently, Qian et al. [18] reported another comprehensive review on the antifouling activities of natural products from marine sources and their synthetic analogs. Another review by Dobretsov et al. [19] reported the progress of biofouling inhibitory activities of marine microorganisms. However, antifouling activities of marine microbes associated with living surfaces are not reviewed comprehensively. The aim of this review was to expand our knowledge on current status of antifouling research from marine microbes associated with macroorganisms.

2. Eco-friendly antifoulants from marine organisms

After the ban of TBT based antifouling paints and environmental concerns associated with other toxic biocides, there is a growing need for the effective eco-friendly antifoulants for marine applications [20,21]. Research interest on natural product antifoulants has been increased in the recent years that was evident from the growing number publications [21,22]. In nature, many marine sessile organisms




Fig. 2. Examples of fouling organisms commonly found on the hard substrata submerged in the marine waters. a: Macroalgae; b: Ascidian; c: Barnacle; d: Bivalves.




Fig. 3. Common antifouling adopted by various industries for biofouling management.

are keeping their surfaces free from fouling organisms [23,24] mainly through the production of secondary metabolites [1,25,26]. The secondary metabolites produced by many marine organisms that showed inhibitory activities against the biofouling organisms would be the ideal lead molecules for the development of natural product antifoulants that can be incorporated into paints [27,28]. The compounds belonging to terpenoids, steroids, carotenoids, phenolics, furanones, alkaloids, peptides and lactones extracted from the marine organisms showed antifouling activities [29]. Among the marine organisms, antifouling activities were largely reported from sponges and corals [24,30,31,32,33,34]. Sponges especially attracted the attention of the researchers due to their close relationship with wide variety of microbes and presence of large number of biologically active secondary metabolites [35]. Another important group attracted the attention of investigators is the ascidians from which good number of antifouling molecules were reported in the literature [36,37]. Antifouling activities were also reported from seaweeds, seagrasses, [38,39], bryozoans [40,41], mangroves and microorganisms [42,43,44,45].

Few antifouling coatings based on natural products from the marine organisms such as Sea Nine- 211, Netsafe and Pearlsafe have already been commercialized [46,47]. During the past five decades bioactive metabolites from the marine environment attracted the attention of researchers all over the world for the discovery of lead compounds in medicine and industry [48,49,50,51,52,53,54,55]. Due to the concern on exploitation of large amount of marine organisms for natural products discovery, marine microbes are considered as the viable source for searching bioactive molecules. Many novel bioactive metabolites with antifouling activities were reported from marine microbes in the literature [56,57,58,59,60].

3. Marine macroorganisms–microbe associations

In the marine environment, macroorganisms are generally colonized by an array of microbial communities from the surrounding waters [61], sometimes in high density up to 40–60% of weight reported for sponge, e.g. Hentchel et al. [62] or diversity (i.e., many strains in an animal, e.g. Li et al. [63]). This association also referred as 'symbiosis' (which includes both ectosymbiosis and endosymbiosis) has been described from all animal and plant groups in the marine realm [64]. The symbionts commonly reported in the literature include microorganisms belonged to bacteria, archaea and unicellular eukaryotes [64]. Generally, the microorganisms attached to marine invertebrates and plants possess more physiological activities than free-living ones [65,66]. The association between invertebrates and microbes occur for many

purposes. For example, these microbes may produce secondary metabolites to enhance their survival in the competitive conditions prevailing on the surface of the host's body [67]. The production of secondary metabolites by these microbes was evident from the studies made by Burkholder et al. [68] in which they reported a bioactive metabolite from the bacterium obtained from the surface of Caribbean seagrass *Thalassia*. Microbes are believed to produce different types of metabolites with pharmacodynamic properties, insecticidal and repellants activities [69,70]. These metabolites are mainly exploited for screening of lead molecules for drugs and other compounds with industrial applications [71,72,73].

The microbes associated with marine organisms and their secondary metabolites may enhance (inductive) or inhibit (non-inductive) the larval settlement of marine organisms. Those microbes which inhibit the larval settlement could be used as a potential source for the exploration of antifouling compounds. Also, bacteria associated with the surface of marine invertebrates are reported to contain a higher proportion of antibacterial and antifouling activities than those occur as planktonic forms [74,75]. This hypothesis was confirmed by the studies made by Long and Azam [76] in which they reported that a major proportion of microbes attached with surfaces produce inhibitory compounds than free-living forms. Several studies indicate that the metabolite believed to be produced by the host organism for the defense purpose is actually originated from the microbes [77,78,79]. To confirm this hypothesis, many investigations were carried out to isolate the bacteria associated with sponges and number of novel metabolites have been reported [80,81]. For example, the cytotoxic macrolide swinholide 1, extracted from the sponge *Theonella swinhoei*, was found to be synthesized by one of the unicellular bacterial symbionts inhabiting the endosome of this sponge species [82]. Hence, microbial associations with higher organisms serve as a sustainable resource for novel biologically active secondary metabolites. This prompted more studies focusing on the metabolites produced by the microorganisms associated with marine macroorganisms.

4. Antifouling activities of microbes associated with marine invertebrates and ascidians

In this review, antifouling activities of bacteria and fungi associated with marine macroorganisms are highlighted with examples (Table 1 and Table 2). Among the marine organisms, microbes associated with sponges and corals topped the list for antifoulant screening assays (Fig. 4). To mention few, Kon-ya et al. [83] isolated upiquinone-8 from a sponge-associated bacterial strain *Alteromonas* sp. which possess

Table 1

Some bacterial strains isolated from the marine macroorganisms with reported antifouling activity/bioactivity.

Host organism	Bacterial strain	Activity	Reference
Sponge	<i>Alteromonas</i> sp.	Antifouling	[83]
Ascidian	<i>Acinetobacter</i> sp.	Antifouling	[84]
Macroalga	<i>Pseudoalteromonas ulvae</i> sp.	Antifouling	[101]
Nudibranchs	<i>Pseudomonas</i> sp.	Antifouling	[94]
Macroalga	<i>Vibrio</i> sp.	Antifouling	[104]
Sponge	<i>Pseudoalteromonas piscicida</i>	Antimicrobial	[138]
Macroalga	<i>Phaeobacter gallaeciensis</i>	Antifouling	[139]
Molluscan	<i>Pseudomonas fulva</i>	Antimicrobial	[140]
Coral	<i>Bacillus horikoshii</i>	Antibacterial/antibiofilm	[141]
Ascidian	<i>Pseudoalteromonas haloplanktis</i>	Antifouling	[118]
Soft coral	<i>Bacillus</i> sp.	Antibacterial	[142]
Seagrass	<i>Bacillus</i> sp.	Antifouling	[107]
Macroalga	<i>Pseudovibrio</i> sp.	Antibacterial	[143]
Sponge	<i>Bacillus licheniformis</i>	Antibacterial/antibiofilm	[144]
Macroalga	<i>Leucothrix mucor</i>	Antifouling	[145]
Macroalga	<i>Streptomyces praecox</i>	Antifouling	[108]
Sponge	<i>B. cereus</i>	Antifouling	[44]
Macroalga	<i>Streptomyces violaceoruber</i>	Antifouling	[110]
Sponge	<i>Bacillus</i> sp.	Antifouling	[90]
Sponge	<i>Bacillus</i> sp.	Antifouling	[89]
Macroalga	<i>Bacillus subtilis</i>	Antibacterial	[146]
Ascidian	<i>P. denitrificans</i>	Antifouling	[103]
Sponge	<i>Pseudomonas fluorescens</i>	Antimicrobial	[147]

inhibitory activities against barnacle larval settlement. Olguin-Urbe et al. [84] isolated an epibiotic bacterium, *Acinetobacter* sp. from the surface of the ascidian *Stomozoa murrayi*. This bacterium produces 6-bromindole-3-carbaldehyde that inhibited the settlement of cyprid of the barnacle *Balanus amphitrite* at a concentrations of 10 mg mL⁻¹. The ecological role of soft coral-associated bacterium *Arthrobacter* sp. against marine biofilm-forming bacteria was highlighted by Radjasa and Sabdono [85]. Another study by Dobretsov and Qian [86] assessed the antifouling effect of epibiotic bacteria isolated from the surface of the soft coral *Dendronephthya* sp. These researchers isolated 11 bacterial strains from the coral surface and found that 2 strains inhibited the settlement of the larva of tubeworm *Hydroides elegans*. Another study by Kanagasabhapathy et al. [87] examined the effects of four strains of Gram positive bacteria (PS2, PS9, PS11 and PS79) isolated from the sponge *Pseudoceratina pupurea* on the growth of bacteria isolated from

Table 2

Examples of fungal strains associated with different macroorganisms. These strains were reported to produce bioactive metabolites with antifouling and antimicrobial activities.

Host organism	Fungal strain	Activity	Reference
Sponge	<i>L. helminthicola</i>	Antifouling	[21]
Gorgonian	<i>Aspergillus</i> sp.	Antibacterial	[148]
Gorgonian	<i>C. lunatus</i>	Antifouling	[88]
Sponge	<i>Aspergillus insuetus</i>	Antifungal, cytotoxic	[149]
Sponge	<i>Aspergillus</i> sp.	Antibacterial, antifouling	[150]
Sponge	<i>Aspergillus</i> sp.	cytotoxic	[151]
Macroalga	<i>Drechslera</i> sp.	Antifouling, antibacterial	[152]
Gorgonian	<i>Penicillium</i> sp.	Antifouling, antibacterial	[22]
Coral	<i>Aspergillus</i>	Antifouling	[153]
Coral	<i>Alternaria</i>	Antifouling	[153]
Gorgonian	<i>Xylariaceae</i> sp.	Antifouling, enzyme-inhibitory activity	[154]
Soft coral	<i>Aspergillus elegans</i>	Antifouling, antibacterial	[155]
Gorgonian	<i>Aspergillus terreus</i>	Antifouling, antiviral	[156]
Sea Anemone	<i>C. lunatus</i>	Antifouling, antifungal	[102]
Sponge	<i>Aspergillus sydowii</i>	Antimicrobial, antiviral	[157]
Gorgonian	<i>Aspergillus</i> sp.	Antimicrobial, antifouling	[158]
Gorgonian	<i>Talaromyces</i> sp.	Cytotoxic, antifouling	[16]
Gorgonian	<i>Scopulariopsis</i> sp.	Antifouling	[91]
Soft coral	<i>Aspergillus unguis</i>	Antifungal	[159]
Gorgonian	<i>P. pinophilum</i>	Antifungal, cytotoxic	[92]

the biofilms and standard strains of genera *Vibrio-Photobacterium*. Two antifouling compounds, 3-methyl-N-(2-phenylethyl) butanamide and cyclo(D-Pro-D-Phe) were isolated from a sponge-associated fungus *Letendreaa helminthicola* [21]. Another fungal strain *Cochliobolus lunatus* associated with the gorgonian *Dichotella gemmacea* collected from the South China Sea showed antifouling activities [88]. The bacterium *Bacillus cereus* isolated from the surface of the sponge *Sigmadocia* sp. was capable of inhibiting the adhesion of biofilm bacteria and microalgae [44]. In a study carried out by Bao et al. [22] screened the antifouling activity of gorgonian derived fungus *Penicillium* sp. SCGAF0023. The symbiotic bacteria associated with sponge *Aplysina gerardogreeni* showed antifouling activities against microfouling organisms such as bacteria and microalgae in laboratory assays [89]. The crude extracts of 52 bacterial strains associated with sponge species were tested for anti-adhesion activities against diatoms by Jin et al. [90] and suggested *Bacillus* sp. as potential source for antifouling compounds. Dihydroquinolin-2(1H)-one containing alkaloids from a fungal strain *Scopulariopsis* sp. associated with the gorgonian coral from the South China Sea showed strong antifouling activities against the larvae of barnacle *Amphibalanus amphitrite* [91]. The compounds produced by a gorgonian derived fungus *Penicillium pinophilum* showed inhibitory activities against the barnacle larvae at non-toxic concentrations [92].

The microbes associated with other macroorganisms were also subjected to extensive studies for antifouling activities. For example, bacterial communities associated with barnacle *B. amphitrite* (= *A. amphitrite*) was found to be active against the settlement of barnacle larvae [93]. The extract of bacterial strain NudMB50-11 isolated from the surface of the nudibranch, *Archidoris pseudoargus* was found to be active against fouling bacteria [4]. Three antibacterial compounds, pyolipic acid, phenazine-1-carboxylic acid and 2-alkylquinol-4-ones extracted from a *Pseudomonas* sp. isolated from the nudibranchs showed strong antifouling activities in both laboratory and field assays [94,95]. Extracellular polymeric substances produced by *Exiguobacterium* sp. associated with the polychaete *Platynereis dumerilii* showed inhibitory activities against the biofilm bacteria [96]. Another study by Shankar et al. [97] evaluated the antibiofilm activities of bacteria associated with polychaetes. The *Acinetobacter* sp. associated with the ascidian *S. murrayi* produces an antifouling metabolite 6-bromindole-3-carbaldehyde that inhibits the settlement of cyprids of the barnacle *B. amphitrite* [84]. *Pseudoalteromonas tunicata*, a bacterial strain isolated from the surface of ascidian *Ciona intestinalis* showed inhibitory activities against larval forms of barnacle, polychaete, ascidian and spores of macroalgae [98,99,100,101]. The resorcylic acid lactones isolated from a sea anemone-associated fungus *C. lunatus* exhibited antifouling activities against the barnacle larvae in laboratory assays [102]. Eight bioactive compounds that belonged to di(1H-indol-3-yl) methane family were isolated from the ascidian associated *Pseudovibrio denitrificans* and all the compounds showed moderate to strong antifouling activities against larval forms of barnacle *B. amphitrite* and bryozoan *Bugula neritina* [103].

5. Antifouling activities of microbes associated with seaweeds and seagrasses 5

Microbes associated with seaweeds and seagrasses were also screened extensively for antifouling activity since they constitute an important source for bioactive substances. Among the seaweed epibiotic bacteria, the genus *Pseudoalteromonas* was highlighted as an important group with antifouling, antimicrobial and cytotoxic activities. For example, *Pseudoalteromonas ulvae* sp. nov., a bacterium with antifouling activities was isolated from the surface of the alga *Ulva lactuca* by Egan et al. [101]. The bacterial strains *Pseudoalteromonas* sp. and *Vibrio alginolyticus* isolated from an alga produce either non-soluble or waterborne metabolites that inhibit larval settlement [104]. Besides, *Pseudoalteromonas* and unidentified bacterial strains




Fig. 4. Examples of marine invertebrate groups commonly reported in the literature for studying microbial symbionts. a: Coral; b: Gorgonian; c: Sponge; d: Soft coral. (Underwater images from the Red Sea).

were also reported to possess antifouling activities. A study by Dobretsov and Qian [105] reported that bacterial strains from the surface of the green alga *Ulva reticulata* showed inhibitive activities against micro- and macrofouling organisms. Active compounds from the cells and culture supernatant of the bacterial strain, FS-55 isolated from the surface of seaweed, *Fucus serratus*, were extracted using solid phase extraction, combined with acrylic based paint resin and showed good antifouling activity against the fouling bacteria [4]. A study carried out by Ma et al. [106] showed the inhibition of common fouling organisms in mariculture by epiphytic bacteria associated with the surface of seaweeds and invertebrates off the Dalian coast in China. The bacterial symbionts of seagrasses *Thalassia hemprichii* and *Enhalus acoroides* were screened for antifouling activities against biofilm-forming bacteria and identified members of the genus *Bacillus* and *Virgibacillus* as active symbionts [107]. The actinomycetes associated with seaweeds and sediments along the coast of Korea were evaluated for their antifouling activity by Cho et al. [108] and reported diketopiperazines as active metabolites. An unidentified epibiotic bacterium from the surface of the seaweed *Sargassum wightii* showed antifouling activity in laboratory and field assays [109]. Two furanone derivatives produced by the bacterium *Streptomyces violaceoruber* isolated from the surface of the brown seaweed *Undaria pinnatifida* showed antifouling activities against macroalgae and mussel larvae [110].

6. *Pseudoalteromonas* associated with marine organisms: A potential genus for antifouling research

Most of the antifouling screening studies on the microbes associated with marine organisms mainly focused on a particular species for detailed study based upon the activities. In general, *Alteromonas*, *Pseudoalteromonas*, *Vibrio* and *Bacillus* were dominant bacterial groups frequently found associated with marine macroorganisms [70,86]. Among these, *Pseudoalteromonas*, a genus reclassified by Gauthier et al. [111] is widely isolated from animal and plant surfaces and reported to produce many bioactive molecules [60,70,100,101,112]. The dominance of this genus on marine living surfaces may be due to the bacteriolytic and algicidal activities [100] which provide effective way for competition with other colonizing organisms [113]. They can also survive under poor nutritional conditions because of the biochemical pathways and production of secondary metabolites which include bioactive compounds and enzymes [114,115].

Pseudoalteromonas received more attention in recent years for natural product research due to the widespread distribution and easy cultivability under laboratory conditions [116]. Previous studies showed that the bioactive molecules present in this genus possess antifouling activities against micro and macrofouling organisms [117,118]. The antifouling activities are mainly related to the presence of yellow and purple pigments in *Pseudoalteromonas* [56] and Franks et al. [119] have identified the yellow pigments as a new member belonging to tambjamine compounds. In addition, many members of this genus are reported to produce extracellular enzymes, toxins and extracellular polymeric substances [100,120,121,122]. The compounds produced by *Pseudoalteromonas* will definitely serve as lead for the development of novel antifoulants. Due to the ecological and biotechnological significance of *Pseudoalteromonas*, more than 50 genomes of this genus have been sequenced [123]. Though, *Pseudoalteromonas* bacteria were isolated from many organisms, given the nature of vast diversity in the marine realm opportunities still exist for concerted research program for searching bioactive molecules with potential antifouling activities from this genus.

7. Advantages of microbes as a source of bioactive metabolites

Majority of the antifouling compounds isolated from the marine organisms are from invertebrates of tropical or subtropical seas where species diversity and resource competition are reported to higher than other ecosystems [50]. Bioactive compounds are synthesized in small quantities by the organisms and occur as a complex mixture [124] and due to that the extraction and purification are labor-intensive and a time-consuming process [124,125]. For the extraction of a bioactive compound from a marine organism, large number of animals or algae would have to be collected from the sea. The collection of large amount of marine organisms particularly sponges; corals and rare species will be a cause of concern from the biodiversity conservation point of view [21]. Contrary to this, if the microbe is considered as a source for the bioactive compound, then the product supply will be ensured by culturing the microorganisms or isolating the genes responsible for the biosynthesis of the particular metabolite [126,127,128,129].

Although the microbes are suggested as an alternative source for marine organisms for antifouling compound discovery, microbial symbionts have complex molecular structures that are hard to

synthesize chemically [21]. The isolation of microbes from the macroorganisms is a bottleneck especially in sponges where the microbe lives inside the tissues. Notwithstanding above issues, the advantages are more when microorganisms associated with marine organisms are used as a source for the exploitation of antifouling compounds. The microbes can be cultured in the laboratory using appropriate culture media and conditions, though there are some limitations to culture some strains using traditional methods. Jensen et al. [130] confirmed higher recovery of cultivable bacteria from marine algal surfaces using culture-dependent methods. The fermentation process allows us to extract good quantity of metabolites from the microbes for bioassays [131]. The microbial strains will grow in laboratory under optimum temperature, pH, and nutrient conditions. Most of the previous studies suggested that microorganisms will grow successfully under conditions that mimic the physical and chemical characters of the natural environment. For example, Okazaki et al. [132] reported that a marine isolate from the shallow sea mud produce antibiotics only when supplemented with powdered *Laminaria* in the growth medium. In addition, there are possibilities that bacterial strains of the same species can produce different bioactive compounds depending on culture conditions and thus increasing the prospective number of valuable compounds [4]. Some of the microorganisms which showed antifouling activities in laboratory assays are failed to exhibit the same in field assays. For the development of natural product based antifouling coatings, there is a need for the evaluation of antifouling potential of the compound through laboratory and field trails (Fig. 5).

The antifouling performance of the bioactive compounds isolated from the marine microbes can be tested in the laboratory against biofilm-forming bacteria, diatoms and barnacle larvae as target organisms. An ideal natural product antifouling compound will act different ways on the target organisms (Fig. 6). Generally, the compound should prevent the formation of biofilms, which is considered as a cue for the further settlement of invertebrate larvae in the marine environment. The extracts or compounds from the microbes isolated from the marine macroorganisms showed inhibitory activities against an array of biofilm-forming bacteria [44,45,97]. The main mechanism of antibiofilm activities of microbial strains associated with marine macroorganisms includes antibiotic activity and anti-adhesion property and affects the extracellular polymers production (EPS) which is essential for biofilm formation [44]. It is

believed that the compounds produced by the microbes associated with marine macroorganisms may exhibit same mode of mechanism (anti-settlement) against larval forms and macroalgal spores though more studies required to confirm the antifouling mechanism.

8. Conclusions and future perspectives

Marine microorganisms are taxonomically diverse and unique, which makes them as potential source for discovery of novel bioactive molecules [133]. In the aquatic ecosystems, microbial communities possess strong affinity towards the living and non-living surfaces [134]. The microbial association with living surfaces in the marine environment provides ample opportunities for bioprospecting natural products. This review clearly confirms the antifouling activities expressed by microbes associated with living surfaces in the marine environment. However, most of the studies were conducted under laboratory conditions and failed to test the compounds in natural water for commercial applications. Hence two possible approaches are suggested for further research. First, as pointed out by Qian et al. [10,18], those metabolites which showed antifouling/antimicrobial activities should be investigated further through different antifouling assays using various target species. These assays may also include field trials by incorporating these metabolites into a suitable paint. Most of the previous investigators used pure compounds or crude extracts from microbes for preparing the antifouling paints [15,94,109]. This approach is rather good than incorporation of surface-associated bacterial strains in suitable paints. Second, more bioprospecting efforts are required to recover novel antifouling molecules from the marine macroorganism–microbe association. Culture dependent methods were previously used to identify the bacterial communities and the advent of molecular methods provides many tools for studying the microbes associated with surfaces [131,135]. Hence, applying the genomic tools along with bioassay guided antifouling assays will yield valid information and novel metabolites from the microbial consortia associated with marine macroorganisms.

The industries certainly request a potent antifouling system with long durability (at least 5 years), cost effective, easy for application and non-toxic to marine ecosystem [136]. However, most of the new antifouling systems failed to meet the above characters. Natural products can be successfully used for antifouling applications by incorporating in a suitable paint. However, preparation of an




Fig. 5. Schematic diagram showing the steps involved in isolation and identification of natural product antifouling compounds from the microbes associated with marine macroorganisms.




Fig. 6. General mode of action of antifouling compounds on fouling organisms (contents adapted from Qian et al. [160]).

antifouling coat using microbial products is a major challenge as these compounds will breakdown rapidly in the environment [137]. Finding a best way to increase the durability of the compound when applied as antifouling coating may provide more opportunities for natural product based antifouling systems. Developing a natural product antifouling coatings based on microbes associated with marine macroorganisms definitely takes much time and efforts. The toxicity of compounds produced by these microbes on non-target organisms in the marine ecosystem also needs to be analyzed before commercial applications [19]. In conclusion, the microbes associated with marine macroorganisms are an untapped source for natural product antifouling compounds and many more novel compounds could be identified through interdisciplinary approach.

Conflict of interest statement

The authors declare that there are no conflict of interest.

References

- [1] Wahl M. Living attached: Aufwuchs, fouling, epibiosis. In: Nagabhushanam R, Thompson M, editors. Fouling organisms of the Indian ocean: Biology and control technology. New Delhi: Oxford and IBH Publishing Company; 1997. p. 31–84.
- [2] Maki J. Biofouling in the marine environment. In: Bitton H, editor. Encyclopedia of environmental microbiology. New York: Wiley & Sons; 2002. p. 610–9.
- [3] Gerhart DJ, Rittschof D, Mayo SW. Chemical ecology and the search for marine antifoulants – Studies of a predator–prey symbiosis. J Chem Ecol 1988;14: 1905–17. <http://dx.doi.org/10.1007/BF01013485>.
- [4] Armstrong E, Boyd KG, Pisacane A, Peppiatt CJ, Burgess JG. Marine microbial natural products in antifouling coatings. Biofouling 2000;16:215–24. <http://dx.doi.org/10.1080/08927010009378446>.
- [5] Qi SH, Zhang S, Yang LH, Qian PY. Antifouling and antibacterial compounds from the gorgonians *Subergorgia suberosa* and *Scrippearia gracillis*. Nat Prod Res 2008; 22:154–66. <http://dx.doi.org/10.1080/14786410701642441>.
- [6] de Nys R, Guenther J. The impact and control of biofouling in marine finfish aquaculture. In: Hellio C, Yebra D, editors. Advances in marine antifouling coatings and technologies. Cambridge (UK): Woodhead Publishing Ltd; 2009. p. 177–221.
- [7] Callow ME, Callow JE. Marine biofouling: A sticky problem. Biologist 2002;49:10–4.
- [8] Global invasive species programme. Marine biofouling: An assessment of risks and management initiatives. Compiled by Jackson L, Global Invasive Species Programme and UNEP Regional Seas Programme. UNEP; 2008[Cited October 25, 2015]. Available from Internet: http://www.issg.org/gisp_publications_reports.htm.
- [9] Hu J, Zhen H, Wan Y, Gao J, An W, An L, et al. Trophic magnification of triphenyltin in a marine food web of Bohai Bay, North China: Comparison to tributyltin. Environ Sci Technol 2006;40:3142–7. <http://dx.doi.org/10.1021/es0514747>.
- [10] Qian PY, Xu Y, Fusetani N. Natural products as antifouling compounds: Recent progress and future perspectives. Biofouling 2009;26:223–34. <http://dx.doi.org/10.1080/08927010903470815>.
- [11] Alzieu C. TBT detrimental effects on oyster culture in France – Evolution since antifouling paint regulation. Oceans 1986;18:1130–4.
- [12] Gibbs PE, Bryan GW. Reproductive failure in populations of the dog-whelk, *Nucella Lapillus*, caused by imposex induced by tributyltin from antifouling paints. J Mar Biol Assoc UK 1986;66:767–77. <http://dx.doi.org/10.1017/S0025315400048414>.
- [13] Swain G. Redefining antifouling coatings. Jpl-Pmc 1999:26–35.
- [14] Hoch M. Organotin compounds in the environment – An overview. Appl Geochem 2001;16:719–43. [http://dx.doi.org/10.1016/S0883-2927\(00\)00067-6](http://dx.doi.org/10.1016/S0883-2927(00)00067-6).
- [15] Yebra DM, Kiil S, Dam-Johansen K. Antifouling technology – Past, present and future steps towards efficient and environmentally friendly antifouling coatings. Prog Org Coat 2004;50:75–104. <http://dx.doi.org/10.1016/j.porgcoat.2003.06.001>.
- [16] Yonehara Y. Recent topics on marine antifouling coatings. Bull Soc Sea Water Sci Jpn 2000;54:7–12. <http://dx.doi.org/10.11457/swsj1965.54.7>.
- [17] Thomas KV. The environmental fate and behavior of antifouling paint booster biocides: A review. Biofouling 2001;17:73–86. <http://dx.doi.org/10.1080/08927010109378466>.

- [18] Qian PY, Li Z, Xu Y, Li Y, Fusetani N. Mini-review: Marine natural products and their synthetic analogs as antifouling compounds: 2009–2014. *Biofouling* 2015;31:101–22. <http://dx.doi.org/10.1080/08927014.2014.997226>.
- [19] Dobretsov S, Abed RMM, Tepilitki M. Mini-review: Inhibition of biofouling by marine microorganisms. *Biofouling* 2013;29:423–41. <http://dx.doi.org/10.1080/08927014.2013.776042>.
- [20] Clare AS. Towards non-toxic antifouling. *J Mar Biotechnol* 1998;6:3–6.
- [21] Yang LH, Miao L, Lee OO, Li X, Xiong H, Pang KL, et al. Effect of culture conditions on antifouling compound production of a sponge-associated fungus. *Appl Microbiol Biotechnol* 2007;74:1221–31. <http://dx.doi.org/10.1007/s00253-006-0780-0>.
- [22] Bao J, Sun YL, Zhang XY, Han Z, Gao HC, He F, et al. Antifouling and antibacterial polyketides from marine gorgonian coral-associated fungus *Penicillium* sp. *SCSAGF* 0023. *J Antibiot* 2013;66:219–23. <http://dx.doi.org/10.1038/ja.2012.110>.
- [23] Keifer PA, Rinehart KL. Renillafoulin, antifouling diterpenes from the sea pansy *Renilla reniformis* (Octocorallia). *J Org Chem* 1986;51:4450–4. <http://dx.doi.org/10.1021/jo00373a020>.
- [24] Yang LH, Lee OO, Jin T, Li XC, Qian PY. Antifouling properties of 10 β -formamidokalihinol-A and kalihinol A isolated from the marine sponge *Acanthella cavernosa*. *Biofouling* 2006;22:23–32. <http://dx.doi.org/10.1080/08927010500498623>.
- [25] Hentschel U, Schmid M, Wagner M, Fieseler L, Gernert C, Hacker J. Isolation and phylogenetic analysis of bacteria with antimicrobial activities from the Mediterranean sponges *Aplysina aerophoba* and *Aplysina cavernicola*. *FEMS Microbiol Ecol* 2001;35:305–12. <http://dx.doi.org/10.1111/j.1574-6941.2001.tb00816.x>.
- [26] Pawlik JR. Antipredatory defensive roles of natural products from marine invertebrates. In: Fattorusso E, Gerwick WH, Tagliatalela-Scafati O, editors. *Handbook of marine natural products*. NY: Springer; 2012. p. 677–710.
- [27] Wahl M, Jensen PR, Fenical W. Chemical control of bacterial epibiosis on ascidians. *Mar Ecol Prog Ser* 1994;110:45–57. <http://dx.doi.org/10.3354/meps110045>.
- [28] Clare AS. Marine natural product antifoulants: Status and potential. *Biofouling* 1996;9:211–29. <http://dx.doi.org/10.1080/08927019609378304>.
- [29] Feng D, Ke C, Li S, Lu C, Guo F. Pyrethroids as promising marine antifoulants: Laboratory and field studies. *Mar Biotechnol* 2009;11:153–60. <http://dx.doi.org/10.1007/s10126-008-9130-9>.
- [30] Sera Y, Adachi K, Nishida F, Shizuri Y. A new sesquiterpene as an antifouling substance from a Palauan marine sponge, *Dysidea herbacea*. *J Nat Prod* 1999;62:395–6. <http://dx.doi.org/10.1021/np980440s>.
- [31] Hedio C, Tsoukatou M, Maréchal J, Aldred N, Beaupoil C, Clare AS, et al. Inhibitory effects of Mediterranean sponge extracts and metabolites on larval settlement of the barnacle *Balanus amphitrite*. *Mar Biotechnol* 2005;7:297–305. <http://dx.doi.org/10.1007/s10126-004-3150-x>.
- [32] Limna Mol VP, Raveendran TV, Abhilash KR, Parameswaran PS. Inhibitory effect of Indian sponge extracts on bacterial strains and larval settlement of the barnacle, *Balanus amphitrite*. *Int Biodeterior Biodegrad* 2010;64:506–10. <http://dx.doi.org/10.1016/j.ibiod.2010.06.003>.
- [33] Ribeiro SM, Rogers R, Rubem AC, Da Gama BP, Muricy G, Pereira RC. Antifouling activity of twelve demosponges from Brazil. *Braz J Biol* 2013;73:501–6. <http://dx.doi.org/10.1590/S1519-69842013000300006>.
- [34] Dobretsov S, Al-Wahaibi ASM, Lai D, Al-Sabahi J, Claerebout M, Proksch P, et al. Inhibition of bacterial fouling by soft coral natural products. *Int Biodeterior Biodegrad* 2015;98:53–8. <http://dx.doi.org/10.1016/j.ibiod.2014.10.019>.
- [35] Taylor MW, Radax R, Steger D, Wagner M. Sponge-associated microorganisms: Evolution, ecology, and biotechnological potential. *Microbiol Mol Biol Rev* 2007;71:295–47. <http://dx.doi.org/10.1128/MMBR.00040-06>.
- [36] Mayzel B, Haber M, Ilan M. Chemical defense against fouling in the solitary ascidian *Phallusia nigra*. *Biol Bull* 2014;227:232–41.
- [37] Trepos R, Cervin G, Hedio C, Pavia H, Stensen W, Stensvåg K, et al. Antifouling compounds from the sub-arctic ascidian *Synoicum pulmonaria*: Synoxazolindinones A and C, pulmonarins A and B, and synthetic analogues. *J Nat Prod* 2014;77:2105–13. <http://dx.doi.org/10.1021/np500503z>.
- [38] De Nys R, Steinberg PD, Willemsen P, Dworjanyan SA, Gabelish CL, King RJ. Broad spectrum effects of secondary metabolites from the red algae *Delisea pulchra* in antifouling assays. *Biofouling* 1995;8:259–71. <http://dx.doi.org/10.1080/08927019509378279>.
- [39] Hedio C, Bergé JP, Beaupoil C, Le Gal Y, Bourgougnon N. Screening of marine algal extracts for anti-settlement activities against microalgae and macroalgae. *Biofouling* 2002;18:205–15. <http://dx.doi.org/10.1080/08927010290010137>.
- [40] Walls JT, Ritz DA, Blackman AJ. Fouling, surface bacteria and antibacterial agents of four bryozoans species found in Tasmania, Australia. *J Exp Mar Biol Ecol* 1993;169:1–13. [http://dx.doi.org/10.1016/0022-0981\(93\)90039-Q](http://dx.doi.org/10.1016/0022-0981(93)90039-Q).
- [41] Kon-ya K, Shimidzu N, Adachi Miki W. 2,5,6-Tribromo-1-methylgramine, an antifouling substance from the marine bryozoans *Zoobrottryon pellucidum*. *Fish Sci* 1994;60:773–5.
- [42] Burgess JG, Jordan EM, Bregu M, Mearns-Spragg A, Boyd KG. Microbial antagonism: A neglected avenue of natural products research. *J Biotechnol* 1999;70:27–32. [http://dx.doi.org/10.1016/S0168-1656\(99\)00054-1](http://dx.doi.org/10.1016/S0168-1656(99)00054-1).
- [43] Kennedy J, Baker P, Piper C, Cotter PD, Walsh M, Mooij MJ, et al. Isolation and analysis of bacteria with antimicrobial activities from the marine sponge *Haliclona simulans* collected from Irish waters. *Mar Biotechnol* 2009;11:384–96. <http://dx.doi.org/10.1007/s10126-008-9154-1>.
- [44] Sathesh S, Soniyamby AR, Sunjaiy Shankar CV, Punitha SMJ. Antifouling activities of marine bacteria associated with the sponge (*Sigmodocia* sp.). *J Ocean Univ China* 2012;11:354–60. <http://dx.doi.org/10.1007/s11802-012-1927-5>.
- [45] Viju N, Anitha A, Sharmin Vini S, Sunjaiy Shankar CV, Sathesh S, Punitha SMJ. Antibiofilm activities of extracellular polymeric substances produced by bacterial symbionts of seaweeds. *Indian J Geo-Mar Sci* 2014;43:1–11.
- [46] Jacobson AH, Willingham GL. Sea-nine antifoulant: An environmentally acceptable alternative to organotin antifoulants. *Sci Total Environ* 2000;258:103–10. [http://dx.doi.org/10.1016/S0048-9697\(00\)00511-8](http://dx.doi.org/10.1016/S0048-9697(00)00511-8).
- [47] De Nys R, Ison O. Evaluation of antifouling products developed for the Australian pearl industry. Fisheries Research and Development Corporation; 2004.
- [48] Cooper EL. Drug discovery, CAM and natural products. *Evid Based Complement Alternat Med* 2004;1:215–7. <http://dx.doi.org/10.1093/ecam/neh032>.
- [49] Jones P, Cottrell MT, Kirchner DL, Dexter SC. Bacterial community structure of biofilms on artificial surfaces in an estuary. *Microb Ecol* 2007;53:153–62. <http://dx.doi.org/10.1007/s00248-006-9154-5>.
- [50] Dunlap WC, Battershill CN, Liprott CH, Cobb RE, Bourne DG, Jaspars M, et al. Biomedicinals from the phytosymbionts of marine invertebrates: A molecular approach. *Methods* 2007;42:358–76. <http://dx.doi.org/10.1016/j.jymeth.2007.03.001>.
- [51] Devi P, Wahidullah S, Rodrigues C, Souza LD. The Sponge-associated bacterium *Bacillus licheniformis* SAB1: A source of antimicrobial compounds. *Mar Drugs* 2010;8:1203–12. <http://dx.doi.org/10.3390/md8041203>.
- [52] Mayer AMS, Rodríguez AD, Berlinck RGS, Fusetani N. Marine pharmacology in 2007–8: Marine compounds with antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiprotozoal, antituberculosis, and antiviral activities; affecting the immune and nervous system, and other miscellaneous mechanisms of action. *Comp Biochem Physiol Part C: Toxicol Pharmacol* 2011;153:191–222. <http://dx.doi.org/10.1016/j.cbpc.2010.08.008>.
- [53] Felczykowska A, Bloch SK, Nejman-Faleńczyk B, Barańska S. Metagenomic approach in the investigation of new bioactive compounds in the marine environment. *Acta Biochim Pol* 2012;59:501–5.
- [54] Cong L, Liang W, Wu Y, Li C, Chang Y, Dong L, et al. High-level soluble expression of the functional peptide derived from the C-terminal domain of the sea cucumber lysozyme and analysis of its antimicrobial activity. *Electron J Biotechnol* 2014;17:280–6. <http://dx.doi.org/10.1016/j.ejbt.2014.09.001>.
- [55] Cortés Y, Hormazábal E, Leal H, Urzúa A, Mutis A, Parra L, et al. Novel antimicrobial activity of a dichloromethane extract obtained from red seaweed *Ceramium rubrum* (Hudson) (Rhodophyta: Florideophyceae) against *Yersinia ruckeri* and *Saprolegnia parasitica*, agents that cause diseases in salmonids. *Electron J Biotechnol* 2014;17:126–31. <http://dx.doi.org/10.1016/j.ejbt.2014.04.005>.
- [56] Egan S, James S, Holmström C, Kjelleberg S. Correlation between pigmentation and antifouling compounds produced by *Pseudoalteromonas tunicata*. *Environ Microbiol* 2002;4:433. <http://dx.doi.org/10.1046/j.1462-2920.2002.00322.x>.
- [57] Bhattarai HD, Ganti VS, Paudel B, Lee YK, Lee HK, Hong YK, et al. Isolation of antifouling compounds from the marine bacterium, *Shewanella oneidensis* SCH0402. *World J Microbiol Biotechnol* 2007;23:243–9. <http://dx.doi.org/10.1007/s11274-006-9220-7>.
- [58] Bowman JP. Bioactive compound synthetic capacity and ecological significance of marine bacterial genus *Pseudoalteromonas*. *Mar Drugs* 2007;5:220–41. <http://dx.doi.org/10.3390/md20070017>.
- [59] Ortega-Morales BO, Chan-Bacab MJ, Miranda-Tello E, Fardeau ML, Carrero JC, Stein T. Antifouling activity of sessile bivalves derived from marine surfaces. *J Ind Microbiol Biotechnol* 2008;35:9–15. <http://dx.doi.org/10.1007/s10295-007-0260-2>.
- [60] Soliev AB, Hosokawa K, Enomoto K. Bioactive pigments from marine bacteria: Applications and physiological roles. *Evid Based Complement Alternat Med* 2011;2011:1–17. <http://dx.doi.org/10.1155/2011/670349>.
- [61] Wahl M, Goecke F, Labes A, Dobretsov S, Weinberger F. The second skin: Ecological role of epibiotic biofilms on marine organisms. *Front Microbiol* 2012;3:1–21. <http://dx.doi.org/10.3389/fmicb.2012.00292>.
- [62] Hentschel U, Usher KM, Taylor MW. Marine sponges as microbial fermenters. *FEMS Microbiol Ecol* 2006;55:167–77. <http://dx.doi.org/10.1111/j.1574-6941.2005.00046.x>.
- [63] Li ZY, He LM, Wu J, Jiang Q. Bacterial community diversity associated with four marine sponges from the South China Sea based on 16S rDNA-DGGE fingerprinting. *J Exp Mar Biol Ecol* 2006;329:75–85. <http://dx.doi.org/10.1016/j.jembe.2005.08.014>.
- [64] Haygood MG, Schmidt EW, Davidson KS, Faulkner JD. Microbial symbionts of marine invertebrates: Opportunities for microbial biotechnology. *J Mol Microbiol Biotechnol* 1999;1:33–43.
- [65] Bonar DB, Weiner RM, Colwell RR. Microbial–invertebrate interactions and potential for biotechnology. *Microb Ecol* 1986;12:101–10. <http://dx.doi.org/10.1007/BF02153225>.
- [66] Okami Y. Marine microorganisms as a source of bioactive agents. *Microb Ecol* 1986;12:65–78. <http://dx.doi.org/10.1007/BF02153223>.
- [67] Zheng L, Han X, Chen H, Lin W, Yan X. Marine bacteria associated with marine macroorganisms: The potential antimicrobial resources. *Ann Microbiol* 2005;55:119–24.
- [68] Burkholder PR, Pfister RM, Leitz FH. Production of a pyrrole antibiotic by a marine bacterium. *Appl Microbiol* 1966;14:649–53.
- [69] Demain AL. Pharmaceutically active secondary metabolites of microorganisms. *Appl Microbiol Biotechnol* 1999;52:455–63. <http://dx.doi.org/10.1007/s002530051546>.
- [70] Chelossi E, Milanese M, Milano A, Pronzato R, Riccardi G. Characterisation and antimicrobial activity of epibiotic bacteria from *Petrosia ficiformis* (Porifera, Demospongiae). *J Exp Mar Biol Ecol* 2004;309:21–33. <http://dx.doi.org/10.1016/j.jembe.2004.03.006>.
- [71] Faulkner DJ. Marine natural products. *Nat Prod Rep* 2000;17:7–55. <http://dx.doi.org/10.1039/a809395d>.
- [72] Huisman G, Gray D. Towards novel processes for the fine-chemical and pharmaceutical industries. *Curr Opin Biotechnol* 2002;13:352–8. [http://dx.doi.org/10.1016/S0958-1669\(02\)00335-X](http://dx.doi.org/10.1016/S0958-1669(02)00335-X).
- [73] Rizzo C, Michaud L, Hörmann B, Gerçe B, Sylødatk C, Hausmann R, et al. Bacteria associated with sabellids (Polychaeta: Annelida) as a novel source of surface active compounds. *Mar Pollut Bull* 2013;70:125–33. <http://dx.doi.org/10.1016/j.marpolbul.2013.02.020>.

- [74] Fenical W. Chemical studies of marine bacteria: Developing a new resource. *Chem Rev* 1993;93:1673–83. <http://dx.doi.org/10.1021/cr00021a001>.
- [75] Dash S, Jin C, Lee OO, Xu Y, Qian PY. Antibacterial and antilarval-settlement potential and metabolite profiles of novel sponge-associated marine bacteria. *J Ind Microb Biotechnol* 2009;36:1047–56. <http://dx.doi.org/10.1007/s10295-009-0588-x>.
- [76] Long RA, Azam F. Antagonistic interactions among marine pelagic bacteria. *Appl Environ Microbiol* 2001;67:4975–83. <http://dx.doi.org/10.1128/AEM.67.11.4975-4983.2001>.
- [77] Simidu U, Noguchi T, Hwang DF, Shida Y, Hashimoto K. Marine bacteria which produce tetrodotoxin. *Appl Environ Microbiol* 1987;53:1714–5.
- [78] Elyakov GB, Kuznetsova T, Mikhailov VV, Maltsev II, Voinov VG, Fedoreyev SA. Brominated diphenyl ethers from a marine bacterium associated with the sponge *Dysidea* sp. *Experientia* 1991;47:632–3. <http://dx.doi.org/10.1007/BF01949894>.
- [79] Kobayashi J, Ishibachi MI. Bioactive metabolites of symbiotic marine microorganisms. *Chem Rev* 1993;93:1753–69. <http://dx.doi.org/10.1021/cr00021a005>.
- [80] Unson MD, Faulkner DJ. Cyanobacterial symbiont biosynthesis of chlorinated metabolites from *Dysidea herbacea* (Porifera). *Experientia* 1993;49:349–53. <http://dx.doi.org/10.1007/BF01923420>.
- [81] Unson MD, Holland ND, Faulkner DJ. A brominated secondary metabolite synthesized by the cyanobacterial symbiont of a marine sponge and accumulation of the crystalline metabolite in the sponge tissue. *Mar Biol* 1994;119:1–11. <http://dx.doi.org/10.1007/BF00350100>.
- [82] Lee YK, Lee JH, Lee HK. Microbial symbiosis in marine sponges. *J Microbiol* 2001;39:254–64.
- [83] Kon-Ya K, Shimidzu N, Otaki N, Yokoyama A, Adachi K, Miki W. Inhibitory effect of bacterial ubiquinones on the settling of barnacle, *Balanus amphitrite*. *Experientia* 1995;51:153–5. <http://dx.doi.org/10.1007/BF01929360>.
- [84] Olguin-Urribe G, Abou-Mansour E, Boulanger A, Debard H, Francisco C, Combaut G. 6-Bromoindole-3-carbaldehyde, from an *Acinetobacter* sp. bacterium associated with the ascidian *Stomozoa murrayi*. *J Chem Ecol* 1997;23:2507–21. <http://dx.doi.org/10.1023/B:JOEC.0000006663.28348.03>.
- [85] Radjasa OK, Sabdono A. Ecological role of a softcoral-associated bacterium *Arthrobacter* sp. on marine biofilm-forming bacteria. *Microbiology* 2008;2:84–8.
- [86] Dobretsov S, Qian PY. The role of epibiotic bacteria from the surface of the soft coral *Dendronephthya* sp. in the inhibition of larval settlement. *J Exp Mar Biol Ecol* 2004;299:35–50. <http://dx.doi.org/10.1016/j.jembe.2003.08.011>.
- [87] Kanagasabhapathy M, Sasaki H, Haldar S, Yamasaki S, Nagata S. Antibacterial activities of marine epibiotic bacteria isolated from brown algae of Japan. *Ann Microbiol* 2006;56:167–73. <http://dx.doi.org/10.1007/BF03175000>.
- [88] Shao CL, Wu HX, Wang CY, Liu QA, Xu Y, Wei MY, et al. Potent antifouling resorcylic acid lactones from the gorgonian-derived fungus *Cochliobolus lunatus*. *J Nat Prod* 2011;74:629–33. <http://dx.doi.org/10.1021/np100641b>.
- [89] Aguila-Ramírez RN, Hernández-Guerrero CJ, González-Acosta B, Id-Daoud G, Hewitt S, Pope J, et al. Antifouling activity of symbiotic bacteria from sponge *Aplysina Gerardogreeni*. *Int Biodeterior Biodegrad* 2014;90:64–70. <http://dx.doi.org/10.1016/j.ibiod.2014.02.003>.
- [90] Jin C, Xin X, Yu S, Qiu J, Miao L, Feng K, et al. Antidiatom activity of marine bacteria associated with sponges from San Juan Island, Washington. *World J Microbiol Biotechnol* 2013;30:1325–34. <http://dx.doi.org/10.1007/s11274-013-1557-0>.
- [91] Shao CH, Xu RF, Wang CY, Qian PY, Wang KL, Wei MY. Potent antifouling marine Dihydroquinolin-2(1H)-one-containing alkaloids from the Gorgonian Coral-derived Fungus *Scopulariopsis* sp. *Mar Biotechnol* 2015;17:408–15. <http://dx.doi.org/10.1007/s10126-015-9628-x>.
- [92] Zhao DL, Shao CL, Zhang Q, Wang KL, Guan FF, Shi T, et al. Azaphilone and diphenyl ether derivatives from a gorgonian-derived strain of the fungus *Penicillium pinophilum*. *J Nat Prod* 2015;78:2310–4. <http://dx.doi.org/10.1021/acs.jnatprod.5b00575>.
- [93] Mary A, Mary V, Rittschof D, Nagabhushanam R. Bacterial-barnacle interaction: Potential of using juncellins and antibiotics to alter structure of bacterial communities. *J Chem Ecol* 1993;19:2155–67. <http://dx.doi.org/10.1007/BF00979654>.
- [94] Burgess JG, Boyd KG, Armstrong E, Jiang Z, Yan L, Berggren M, et al. The development of a marine natural product-based antifouling paint. *Biofouling* 2003;19:197–205. <http://dx.doi.org/10.1080/0892701031000061778>.
- [95] Eguía E, Trueba A. Application of marine biotechnology in the production of natural biocides for testing on environmentally innocuous antifouling coatings. *J Coat Technol Res* 2007;4:191–202. <http://dx.doi.org/10.1007/s11998-007-9022-3>.
- [96] Rajasree V, Sunjaiy Shankar CV, Satheesh S, Punitha SPJ. Biofilm inhibitory activity of extracellular polymeric substance produced by *Exiguobacterium* sp. associated with the polychaete *Platynereis dumerilii*. *Thalassus* 2014;30:13–9.
- [97] Shankar CVS, Satheesh S, Viju N, Punitha SMJ. Antibacterial and biofilm inhibitory activities of bacteria associated with polychaetes, 3; 2015 495–502. <http://dx.doi.org/10.12980/JCLM.3.2015JCLM-2015-0012>.
- [98] Holmström C, Egan S, Franks A, McCloy S, Kjelleberg S. Antifouling activities expressed by marine surface associated *Pseudoalteromonas* species. *FEMS Microbiol Ecol* 2002;41:47–58. <http://dx.doi.org/10.1111/j.1574-6941.2002.tb00965.x>.
- [99] James SG, Holmström C, Kjelleberg S. Purification and characterization of a novel antibacterial protein from the marine bacterium D2. *Appl Environ Microbiol* 1996;62:2783–8.
- [100] Holmström C, Kjelleberg S. Marine *Pseudoalteromonas* species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiol Ecol* 1999;30:285–93. <http://dx.doi.org/10.1111/j.1574-6941.1999.tb00656.x>.
- [101] Egan S, Holmström C, Kjelleberg S. *Pseudoalteromonas ulvae* sp. nov., a bacterium with antifouling activities isolated from the surface of a marine alga. *Int J Syst Evol Microbiol* 2001;51:1499–504. <http://dx.doi.org/10.1099/00207713-51-4-1499>.
- [102] Liu QA, Shao CL, Gu YC, Blum M, Gan LS, Wang KL, et al. Antifouling and fungicidal resorcylic acid lactones from the sea anemone-derived fungus *Cochliobolus lunatus*. *J Agric Food Chem* 2014;62:3183–91. <http://dx.doi.org/10.1021/jf500248z>.
- [103] Wang KL, Xu Y, Lu L, Li Y, Han Z, Zhang J, et al. Low-toxicity diindol-3-ylmethanes as potent antifouling compounds. *Mar Biotechnol* 2015;17:624–32. <http://dx.doi.org/10.1007/s10126-015-9656-6>.
- [104] Harder T, Dobretsov S, Qian PY. Waterborne polar macromolecules act as algal antifoulants in the seaweed *Ulva reticulata*. *Mar Ecol Prog Ser* 2004;274:133–41. <http://dx.doi.org/10.3354/meps274133>.
- [105] Dobretsov S, Qian PY. Effect of bacteria associated with the green alga *Ulva reticulata* on marine micro- and macrofouling. *Biofouling* 2002;18:217–28. <http://dx.doi.org/10.1080/08927010290013026>.
- [106] Ma Y, Liu P, Yu S, Li D, Cao S. Inhibition of common fouling organisms in mariculture by epiphytic bacteria from the surfaces of seaweeds and invertebrates. *Acta Ecol Sin* 2009;29:222–6. <http://dx.doi.org/10.1016/j.chnaes.2009.08.004>.
- [107] Marhaeni B, Radjasa OK, Khoeri MM, Sabdono A, Bengen DG, Sudoyo H. Antifouling activity of bacterial symbionts of seagrasses against marine biofilm-forming bacteria. *J Environ Prot* 2011;2:1245–9. <http://dx.doi.org/10.4236/jep.2011.29143>.
- [108] Cho JY, Kang JY, Hong YK, Baek HH, Shin HW, Kim MS. Isolation and structural determination of the antifouling diketopiperazines from marine-derived *Streptomyces praecox* 291-11. *Biosci Biotechnol Biochem* 2012;76:1116–21. <http://dx.doi.org/10.1271/bbb.110943>.
- [109] Rajasree V, Satheesh S, Vincent SGP. Antifouling activity of marine epibiotic bacterium from the seaweed *Sargassum wightii*. *Thalassus* 2012;28:37–44.
- [110] Hong YK, Cho JY. Effect of seaweed epibiotic bacterium *Streptomyces violaceoruber* SCH-09 on marine fouling organisms. *Fish Sci* 2013;79:469–75. <http://dx.doi.org/10.1007/s12562-013-0604-y>.
- [111] Gauthier G, Gauthier M, Christen R. Phylogenetic analysis of the genera *Alteromonas*, *Shewanella*, and *Moritella* using genes coding for small-subunit rRNA sequences and division of the genus *Alteromonas* into two genera, *Alteromonas* (emended) and *Pseudoalteromonas* gen. nov., and proposal of twelve new species combinations. *Int J Syst Bacteriol* 1995;45:755–61. <http://dx.doi.org/10.1099/00207713-45-4-755>.
- [112] De Rosa S, Milone A, Kujumgiev A, Stefanov K, Nechev I, Popov S. Metabolites from a marine bacterium *Pseudomonas/Alteromonas*, associated with the sponge *Dysidea fragilis*. *Comp Biochem Physiol B Biochem Mol Biol* 2000;126:391–6. [http://dx.doi.org/10.1016/S0305-0491\(00\)00208-X](http://dx.doi.org/10.1016/S0305-0491(00)00208-X).
- [113] Pukall R, Kramer I, Rohde M, Stackebrandt E. Microbial diversity of cultivatable bacteria associated with the North Sea bryozoan *Fuflustra foliacea*. *Syst Appl Microbiol* 2001;24:623–33. <http://dx.doi.org/10.1078/0723-2020-00073>.
- [114] Ivanova EP, Bakunina IY, Nedashkovskaya OI, Gorshkova NM, Alexeeva YV, Zelepuga EA, et al. Ecophysiological variabilities in ectohydrolytic enzyme activities of some *Pseudoalteromonas* species, *P. citrea*, *P. issachenkoi*, and *P. nigrifaciens*. *Curr Microbiol* 2003;46:6–10. <http://dx.doi.org/10.1007/s00284-002-3794-6>.
- [115] Al Khudary R, Stösser NI, Qoura F, Antranikian G. *Pseudoalteromonas arctica* sp. nov., an aerobic, psychrotolerant, marine bacterium isolated from Spitzbergen. *Int J Syst Evol Microbiol* 2008;58:2018–24. <http://dx.doi.org/10.1099/ijs.0.64963-0>.
- [116] Ivanova EP, Gorshkova NM, Zhukova NV, Lysenko AM, Zelepuga EA, Prokofeva NG, et al. Characterization of *Pseudoalteromonas distincta*-like sea-water isolates and description of *Pseudoalteromonas aliena* sp. nov. *Int J Syst Evol Microbiol* 2004;54:1431–7. <http://dx.doi.org/10.1099/ijs.0.03053-0>.
- [117] Holmström C, James S, Neilan BA, White DC, Kjelleberg S. *Pseudoalteromonas tunicata* sp. nov., a bacterium that produces antifouling agents. *Int J Syst Bacteriol* 1998;48:1205–12. <http://dx.doi.org/10.1099/00207713-48-4-1205>.
- [118] Ma Y, Liu P, Zhang Y, Cao S, Li D, Chen W. Inhibition of spore germination of *Ulva pertusa* by the marine bacterium *Pseudoalteromonas haloplanktis* C14. *Acta Oceanol Sin* 2010;29:69–78. <http://dx.doi.org/10.1007/s13131-010-0009-z>.
- [119] Franks A, Haywood P, Holmström C, Egan S, Kjelleberg S, Kumar N. Isolation and structure elucidation of a novel yellow pigment from the marine bacterium *Pseudoalteromonas tunicata*. *Molecules* 2005;10:1286–91. <http://dx.doi.org/10.3390/10101286>.
- [120] Bowman JP. *Pseudoalteromonas prydzensis* sp. nov., a psychrotrophic, halotolerant bacterium from Antarctic sea ice. *Int J Syst Bacteriol* 1998;48:1037–41.
- [121] Ivanova EP, Kiprianova EA, Mikhailov VV, Levanova GF, Garagulya AD, Gorshkova NM, et al. Phenotypic diversity of *Pseudoalteromonas citrea* from different marine habitats and emendation of the description. *Int J Syst Evol Microbiol* 1998;48:247–56. <http://dx.doi.org/10.1099/00207713-48-1-247>.
- [122] Skovhus TL, Holmström C, Kjelleberg S, Dahllöf I. Molecular investigation of the distribution, abundance and diversity of the genus *Pseudoalteromonas* in marine samples. *FEMS Microbiol Ecol* 2007;61:348–61. <http://dx.doi.org/10.1111/j.1574-6941.2007.00339.x>.
- [123] Wang P, Yu Z, Li B, Cai X, Zeng Z, Chen X, et al. Development of an efficient conjugation-based genetic manipulation system for *Pseudoalteromonas*. *Microb Cell Fact* 2015;14:11. <http://dx.doi.org/10.1186/s12934-015-0194-8>.
- [124] Gil-Chavez GJ, Villa JA, Ayala-Zavala JF, Basilio Heredia J, Sepulveda D, Yahia EM, et al. Technologies for extraction and production of bioactive compounds to be used as nutraceuticals and food ingredients: An overview. *Compr Rev Food Sci Food Saf* 2013;12:5–23. <http://dx.doi.org/10.1111/1541-4337.12005>.
- [125] Lam KS. Discovery of novel metabolites from marine actinomycetes. *Curr Opin Microbiol* 2006;9:245–51. <http://dx.doi.org/10.1016/j.mib.2006.03.004>.
- [126] Piel J. Bacterial symbionts: Prospects for the sustainable production of invertebrate-derived pharmaceuticals. *Curr Med Chem* 2006;13:39–50. <http://dx.doi.org/10.2174/092986706775197944>.
- [127] Salomon CE, Magarvey NA, Sherman DH. Merging the potential of microbial genetics with biological and chemical diversity: An even brighter future for marine natural product drug discovery. *Nat Prod Rep* 2004;21:105–21. <http://dx.doi.org/10.1039/b301384g>.
- [128] Schmidt EW. Trading molecules and tracking targets in symbiotic interactions. *Nat Chem Biol* 2008;4:466–73. <http://dx.doi.org/10.1038/nchembio.101>.

- [129] Hochmuth T, Piel J. Polyketide synthases of bacterial symbionts in sponges – Evolution-based applications in natural products research. *Phytochemistry* 2009; 70:1841–9. <http://dx.doi.org/10.1016/j.phytochem.2009.04.010>.
- [130] Jensen PR, Kauffman CA, Fenical W. High recovery of culturable bacteria from the surfaces of marine algae. *Mar Biol* 1996;126:1–7. <http://dx.doi.org/10.1007/BF00571371>.
- [131] Penesyan A, Kjelleberg S, Egan S. Development of novel drugs from marine surface associated microorganisms. *Mar Drugs* 2010;8:438–59. <http://dx.doi.org/10.3390/md8030438>.
- [132] Okazaki T, Kitahara T, Okami Y. Studies on marine microorganisms. IV. A new antibiotic SS-228 Y produced by *Chaimia* isolated from shallow sea mud. *J Antibiot* 1975;28:176–84.
- [133] Debbab A, Aly AH, Lin WH, Proksch P. Bioactive compounds from marine bacteria and fungi. *Microb Biotechnol* 2010;3:544–63. <http://dx.doi.org/10.1111/j.1751-7915.2010.00179.x>.
- [134] El Bour M, Ismail-Ben Ali A, Ktari L. Seaweeds epibionts: Biodiversity and potential bioactivities. In: Méndez-Vilas A, editor. *Microbial pathogens and strategies for combating them: Science, technology and education*. Spain: Formatex Research Center; 2013. p. 1298–306 [www.formatex.info/microbiology4/vol2/1298-1306.pdf].
- [135] Ceh J, Raina JB, Soo RM, van Keulen M, Bourne DG. Coral-bacterial communities before and after a coral mass spawning event on Ningaloo Reef. *PLoS One* 2012; 7:3–10. <http://dx.doi.org/10.1371/journal.pone.0036920>.
- [136] Ralston E, Swain G. Bioinspiration—the solution for biofouling control? *Bioinspir Biomim* 2009;4:1–9. <http://dx.doi.org/10.1088/1748-3182/4/1/015007>.
- [137] Helliö C. The potential of marine biotechnology for the development of new antifouling solutions. *J Sci Haliéutiques Aquat* 2010;2:35–4.
- [138] Zheng L, Chen H, Han X, Lin W, Yan X. Antimicrobial screening and active compound isolation from marine bacterium NJ6-3-1 associated with the sponge *Hymeniacidon perleve*. *World J Microbiol Biotechnol* 2005;21:201–6. <http://dx.doi.org/10.1007/s11274-004-3318-6>.
- [139] Rao D, Webb JS, Holmstrom C, Case R, Low A, Steinberg P, et al. Low densities of epiphytic bacteria from the marine alga *Ulva australis* inhibit settlement of fouling organisms. *Appl Environ Microbiol* 2007;73:7844–52. <http://dx.doi.org/10.1128/AEM.01543-07>.
- [140] Romanenko LA, Uchino M, Kalinovskaya NI, Mikhailov VV. Isolation, phylogenetic analysis and screening of marine mollusc-associated bacteria for antimicrobial, hemolytic and surface activities. *Microbiol Res* 2008;163:633–44. <http://dx.doi.org/10.1016/j.micres.2006.10.001>.
- [141] Thenmozhi R, Nithyanand P, Rathna J, Karutha Pandian S. Antibiofilm activity of coral-associated bacteria against different clinical *M* serotypes of *Streptococcus pyogenes*. *FEMS Immunol Med Microbiol* 2009;57:284–94. <http://dx.doi.org/10.1111/j.1574-695X.2009.00613.x>.
- [142] Sulistiyani S, Nugraheni SA, Radjasa OK, Sabdono A, Khoeri MM. Antibacterial activities of bacterial symbionts of soft coral *Sinularia* sp. against tuberculosis bacteria. *J Coast Dev* 2010;14:45–50.
- [143] Penesyan A, Tebben J, Lee M, Thomas T, Kjelleberg S, Harder T, et al. Identification of the antibacterial compound produced by the marine epiphytic bacterium *Pseudovibrio* sp. D323 and related sponge-associated bacteria. *Mar Drugs* 2011;9: 1391–402. <http://dx.doi.org/10.3390/md9081391>.
- [144] Sayem SA, Manzo E, Ciavatta L, Tramice A, Cordone A, Zanfardino A, et al. Antibiofilm activity of an exopolysaccharide from a sponge-associated strain of *Bacillus licheniformis*. *Microb Cell Fact* 2011;10:74. <http://dx.doi.org/10.1186/1475-2859-10-74>.
- [145] Cho JY. Antifouling steroids isolated from red alga epiphyte filamentous bacterium *Leucothrix mucor*. *Fish Sci* 2012;78:683–9. <http://dx.doi.org/10.1007/s12562-012-0490-8>.
- [146] Susilowati R, Sabdono A, Widowati I. Isolation and characterization of bacteria associated with brown algae *Sargassum* spp. from Panjang Island and their antibacterial activities. *Procedia Environ Sci* 2015;23:240–6. <http://dx.doi.org/10.1016/j.proenv.2015.01.036>.
- [147] Wei MY, Wang CY, Liu QA, Shao CL, She ZG, Lin YC. Five sesquiterpenoids from a marine-derived fungus *Aspergillus* sp. Isolated from a Gorgonian *Dichotella gemmacea*. *Mar Drugs* 2010;8:941–9. <http://dx.doi.org/10.3390/md8040941>.
- [148] Cohen E, Koch L, Thu KM, Rahamim Y, Aluma Y, Ilan M, et al. Novel terpenoids of the fungus *Aspergillus insuetus* isolated from the Mediterranean sponge *Psammocinia* sp. collected along the coast of Israel. *Bioorg Med Chem* 2011;19: 6587–93. <http://dx.doi.org/10.1016/j.bmc.2011.05.045>.
- [149] Li D, Xu Y, Shao CL, Yang RY, Zheng CJ, Chen YY, et al. Antibacterial bisabolane-type sesquiterpenoids from the sponge-derived fungus *Aspergillus* sp. *Mar Drugs* 2012; 10:234–41. <http://dx.doi.org/10.3390/md10010234>.
- [150] Sun LL, Shao CL, Chen JF, Guo ZY, Fu XM, Chen M, et al. New bisabolane sesquiterpenoids from a marine-derived fungus *Aspergillus* sp. isolated from the sponge *Xestospongia testudinaria*. *Bioorg Med Chem Lett* 2012;22:1326–9. <http://dx.doi.org/10.1016/j.bmcl.2011.12.083>.
- [151] Abdel-Lateff A, Okino T, Alarif WM, Al-Lihaibi SS. Sesquiterpenes from the marine algalic fungus *Drechslera* sp. *J Saudi Chem Soc* 2013;17:161–5. <http://dx.doi.org/10.1016/j.jscs.2011.03.002>.
- [152] Li YX, Wu HX, Xu Y, Shao CL, Wang CY, Qian PY. Antifouling activity of secondary metabolites isolated from Chinese marine organisms. *Mar Biotechnol* 2013;15: 552–8. <http://dx.doi.org/10.1007/s10126-013-9502-7>.
- [153] Nong XH, Wang YF, Zhang XY, Zhou MP, Xu XY, Qi SH. Termitrem and butyrolactone derivatives from a marine-derived fungus *Aspergillus terreus*. *Mar Drugs* 2014;12: 6113–24. <http://dx.doi.org/10.3390/md12126113>.
- [154] Zheng CJ, Shao CL, Wu LY, Chen M, Wang KL, Zhao DL, et al. Bioactive phenylalanine derivatives and cytochalasins from the soft coral-derived fungus, *Aspergillus elegans*. *Mar Drugs* 2013;11:2054–68. <http://dx.doi.org/10.3390/md11062054>.
- [155] Nong XH, Zheng ZH, Zhang XY, Lu XH, Qi SH. Polyketides from a marine-derived fungus *Xylariaceae* sp. *Mar Drugs* 2013;11:1718–27. <http://dx.doi.org/10.3390/md11051718>.
- [156] Wang JF, Lin XP, Qin C, Liao SR, Wan JT, Zhang TY, et al. Antimicrobial and antiviral sesquiterpenoids from sponge-associated fungus, *Aspergillus sydowii* ZSDS1-F6. *J Antibiot* 2014;67:581–3. <http://dx.doi.org/10.1038/ja.2014.39>.
- [157] Chen M, Wang K, Liu M, She Z, Wang CY. Bioactive steroid derivatives and butyrolactone derivatives from a Gorgonian-derived *Aspergillus* sp. fungus. *Chem Biodivers* 2015;12:1398–406. <http://dx.doi.org/10.1002/cbdv.201400321>.
- [158] Putri DA, Radjasa OK, Pringgenies D. Effectiveness of marine fungal symbiont isolated from soft coral *Sinularia* sp. from Panjang Island as antifungal. *Procedia Environ Sci* 2015;23:351–7. <http://dx.doi.org/10.1016/j.proenv.2015.01.051>.
- [159] Santos OCS, Soares AR, Machado FLS, Romanos MTV, Muricy G, Giambiagi-Marval M, et al. Investigation of biotechnological potential of sponge-associated bacteria collected in Brazilian coast. *Lett Appl Microbiol* 2015;60:140–7. <http://dx.doi.org/10.1111/lam.12347>.
- [160] Qian PY, Chen L, Xu Y. Mini-review: Molecular mechanisms of antifouling compounds. *Biofouling* 2013;29:318–400. <http://dx.doi.org/10.1080/08927014.2013.776546>.