



## Research article

# Increasing pinosylvin production in *Escherichia coli* by reducing the expression level of the gene *fabI*-encoded enoyl-acyl carrier protein reductase



Caheri Salas-Navarrete <sup>a</sup>, Georgina Hernández-Chávez <sup>a</sup>, Noemí Flores <sup>a</sup>, Luz María Martínez <sup>a</sup>, Alfredo Martínez <sup>a</sup>, Francisco Bolívar <sup>a</sup>, Francisco Barona-Gomez <sup>b</sup>, Guillermo Gosset <sup>a,\*</sup>

<sup>a</sup> Instituto de Biotecnología, Departamento de Ingeniería Celular y Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico

<sup>b</sup> Laboratorio Nacional de Genómica para la Biodiversidad, Cinvestav – IPN, Irapuato, Guanajuato, Mexico

## ARTICLE INFO

## Article history:

Received 14 December 2017

Accepted 5 March 2018

Available online 8 March 2018

## Keywords:

4-Coumaroyl-CoA ligase

Enoyl-acyl carrier protein reductase

*Escherichia coli*

Fatty acid synthesis

Metabolic engineering

Natural products

Pinosylvin

Plant secondary metabolites

Recombineering

Stilbene synthase

Stilbenes

## ABSTRACT

**Background:** The plant secondary metabolite pinosylvin is a polyphenol from the stilbene family, which have positive effects on human health. Biotechnological production is an attractive alternative for obtaining this stilbene. In *Escherichia coli*, malonyl-CoA is the precursor for both stilbene and fatty acid syntheses. In this study, with the aim of increasing pinosylvin production, we evaluated a novel approach that is based on reducing the expression of the gene *fabI*, which encodes the enzyme enoyl-acyl carrier protein reductase that is involved in fatty acid synthesis. **Results:** A recombineering method was employed to eliminate the chromosomal -35 promoter sequence and the upstream region of the gene *fabI* in *E. coli* strain W3110. Analysis, employing RT-qPCR, showed that such modification caused a 60% reduction in the *fabI* transcript level in the mutant strain W3110Δ-35*fabI*::Cm compared to the wild type W3110. Synthetic genes encoding a mutant version of 4-coumaroyl-CoA ligase from *Streptomyces coelicolor* A3 with improved catalytic activity employing cinnamic acid as substrate and a stilbene synthase from *Vitis vinifera* were cloned to generate the plasmid pTrc-Sc4CL(M)-VvSTS. The production performance of strains W3110Δ-35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS and W3110/pTrc-Sc4CL(M)-VvSTS was determined in shake flask cultures with Luria-Bertani medium supplemented with 10 g/L glycerol and 3 mM cinnamic acid. Under these conditions, the strain W3110Δ-35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS produced 52.67 mg/L pinosylvin, a level 1.5-fold higher than that observed with W3110/pTrc-Sc4CL(M)-VvSTS. **Conclusion:** A reduction in the transcript level of *fabI* caused by the elimination of the -35 and upstream promoter sequences is a successful strategy to improve pinosylvin production in *E. coli*.

**How to cite:** Salas-Navarrete C, Hernández-Chávez G, Flores N, et al. Increasing pinosylvin production in *Escherichia coli* by reducing expression level of gene *fabI*-encoded enoyl-acyl carrier protein reductase. Electron J Biotechnol 2018;33:11–16. <https://doi.org/10.1016/j.ejbt.2018.03.001>.

© 2018 Pontificia Universidad Católica de Valparaíso. Production and hosting by Elsevier B.V. All rights reserved. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The stilbenes belong to the phenylpropanoid family of natural products, which includes compounds that have positive effects on human health [1,2,3]. The stilbene pinosylvin (3,5-dihydroxy-trans-stilbene) is a low-molecular-weight plant secondary metabolite involved in defense against infectious agents [4]. It has been reported that pinosylvin can protect animal cells against oxidative stress [5]. This effect has been observed in human retinal cells, where pinosylvin applied at a concentration of 5 to 10 μM improved survival after


oxidative stress [6]. It has also been reported that this stilbene can prevent necrosis in bovine aortic endothelial cells [7].

Pinosylvin can be extracted from plant tissues; however, its low concentration (1 to 40 mg/g of pine wood) and the existence of similar compounds complicate its purification [8]. Therefore, an alternative approach for obtaining this stilbene is based on its production utilizing recombinant microorganisms. The pinosylvin biosynthetic pathway starts with L-phenylalanine, which is transformed by the enzyme L-phenylalanine ammonia lyase (PAL) to cinnamic acid. In a subsequent step, the enzyme 4-coumarate/cinnamate:coenzyme A ligase (4CL) converts the cinnamic acid to cinnamoyl-CoA. In the final step, the enzyme stilbene synthase (STS) catalyzes the stepwise condensation of three molecules of malonyl-CoA with one molecule of cinnamoyl-CoA, yielding one molecule of pinosylvin (Fig. 1).

\* Corresponding author.

E-mail address: [gosset@ibt.unam.mx](mailto:gosset@ibt.unam.mx) (G. Gosset).

Peer review under responsibility of Pontificia Universidad Católica de Valparaíso.



**Fig. 1.** Pinosylvin biosynthetic pathway. PAL, phenylalanine ammonia lyase; 4CL, cinnamate 4-hydroxylase; STS, stilbene synthase; ACC, acetyl-CoA carboxylase; ACC, acetyl-CoA carboxylase; PEP, phosphoenolpyruvate; E4P, erythrose-4-phosphate; DHAP, dihydroxyacetone phosphate; PYR, pyruvate; AcCoA, acetyl-CoA; SHK, shikimate; CHA, chorismate; *fabD*, gene encoding malonyl-CoA:acyl carrier protein (ACP) transacylase; *fabB*,  $\beta$ -ketoacyl-ACP synthase I; *fabF*, gene encoding  $\beta$ -ketoacyl-ACP synthase II; *fabH*, gene encoding  $\beta$ -ketoacyl-ACP synthase III; *fabG*, gene encoding NADPH-dependent  $\beta$ -ketoacyl-ACP reductase; *fabZ*, gene encoding  $\beta$ -hydroxyacyl-ACP dehydratase.

Approaches for generating microbial strains with the capacity of synthesizing pinosylvin are based on the expression of the complete or partial pinosylvin biosynthetic pathway, as well as other modifications that increase the availability of precursor molecules. A strain of *Streptomyces venezuelae* was generated for the production of stilbenes by expressing the gene encoding 4CL from *Streptomyces coelicolor* and a codon-optimized gene that codes for an STS from *Arachis hypogaea*. In cultures grown in a complex medium supplemented with 1.2 mM cinnamic acid, 0.6 mg/L of pinosylvin was produced [9].

A synthetic pinosylvin pathway was generated and evaluated in *Escherichia coli* by employing PAL, 4CL, and STS enzymes from several sources. The most suitable pathway configuration was found to consist of genes encoding PAL from *Petroselinum crispum*, 4CL from *S. coelicolor*, and STS from *Pinus strobus*. Directed protein evolution of *P. strobus* STS was performed to obtain two mutants that caused increased pinosylvin production when expressed with the rest of the biosynthetic pathway. An *E. coli* strain expressing such optimized pathway produced 70 mg/L of pinosylvin in the presence of cerulenin. When 3 mg/L of the precursor L-Phe was added to the culture medium, the pinosylvin titer increased to 91 mg/L [10]. Cerulenin is an inhibitor of fatty acid biosynthesis and reduces the consumption of malonyl-CoA by this pathway, thereby making this precursor available for stilbene biosynthesis [11]. However, the high cost of cerulenin

(>\$20/1 mg) would be a major drawback when considering employing it for industrial production.

In another report, a combinatorial approach was applied to generate *E. coli* strains that produce pinosylvin from glycerol and cinnamic acid. The genes encoding 4CL from *Arabidopsis thaliana* and STS from *Vitis vinifera* were expressed in a combinatorial fashion by employing several different ribosome binding site (RBS) sequences and plasmid origins of replication. Once the best combination of RBS and the origin of replication for pinosylvin production was identified, it was determined that *E. coli* Dh5 $\alpha$  was the best producer among seven evaluated strains. To improve strain performance, a strategy to increase malonyl-CoA was evaluated. It consisted on employing clustered regularly interspaced short palindromic repeats interference (CRISPRi) to repress the expression of the genes *fabB*, *fabD*, and *fabF* encoding enzymes from the fatty acid biosynthesis pathway. It was determined that CRISPRi repression of *fabB*, *fabD*, and *fabF* resulted in 1.5-, 1.9-, and 0-fold increased production of pinosylvin, respectively. By employing a strain that has the inducible *fabD* repression, it was possible to transform 0.5 mM cinnamic acid to 47.5 mg/L (0.22 mM) pinosylvin [12].

As shown in the above examples, reduction in the consumption of malonyl-CoA by repression/inhibition of genes and enzymes from the fatty acid biosynthetic pathway results in increased stilbene production. Other approaches aimed at increasing malonyl-CoA availability have

been successfully applied to the production of compounds from the phenylpropanoid family. The malonate assimilation pathway from *Rhizobium trifolii* is composed of a malonyl-CoA synthetase and a malonate transport protein (*matB* and *matC*). Expression of these two genes in *E. coli* has been shown to increase (2S)-pinocembrin production when malonate is supplemented in the medium [13]. The enzyme acetyl-CoA carboxylase (*Acc*) catalyzes the synthesis of malonyl-CoA from acetyl-CoA. Overexpression of this enzyme from *Corynebacterium glutamicum* in *E. coli* from the strong T7 promoter in cultures performed at 25°C resulted in a threefold increase in malonyl-CoA concentration. To increase acetyl-CoA availability for malonyl-CoA synthesis, overexpression of acetyl-CoA synthetase (*Acs*) was combined with deletion of the genes *ackA-pta* and *adhE*, which encode acetate kinase, phosphotransacetylase and alcohol/aldehyde dehydrogenase. When tested in a strain that also overexpresses *Acc* from *C. glutamicum*, it was determined that malonyl-CoA concentration increased 15.7-fold [14].

Various approaches have been evaluated to generate and improve pinosylvin production strains, including increasing malonyl-CoA availability. Even though these strategies have a positive effect regarding the increase in precursor availability, such modifications can have negative effects on strain growth capacity or process cost. For this reason, it is desirable to explore alternative approaches that improve pinosylvin production without compromising strain physiology.

The gene *fabI* encodes enoyl-acyl carrier protein reductase, an enzyme that has a key role in fatty acid biosynthesis. To study the role of this enzyme on malonyl-CoA metabolism, an *E. coli* strain that has a thermosensitive *FabI* mutant has been characterized. It was determined that growth of this strain at a nonpermissive temperature of 42°C resulted in a 30% increase in the malonyl-CoA pool [15]. It should also be noted that *fabI* is an essential gene; therefore, it cannot be deleted in *E. coli*. Based on this information, we decided to explore an approach based on reducing the amount of enoyl-acyl carrier protein reductase to increase the malonyl-CoA pool. The strategy consisted on deleting the -35 promoter sequence of gene *fabI*, expecting a reduction in its transcript level. This strategy would avoid the undesired stress responses caused by increasing the culture temperature to 42°C, if a thermosensitive *FabI* mutant was used [16].

In this work, we constructed an *E. coli* strain that can convert cinnamic acid to pinosylvin and showed the feasibility of reducing the expression level of the gene *fabI*, encoding the enzyme enoyl-acyl carrier protein reductase, by employing a recombinering strategy to delete the -35 and upstream promoter regions. An engineered strain with such modification was found to produce 52.67 mg/L pinosylvin, a level 1.5-fold higher compared to an isogenic wild type strain without reduction in the expression level of *fabI*.

## 2. Materials and methods

### 2.1. Bacterial strains and plasmids

The *E. coli* strain W3110 was employed in this study [17] (Table 1). The genes 4CL(M) from *S. coelicolor* A3 (2) (Sc4CL(M)) and STS from *V. vinifera* (VvSTS) were synthesized with codon optimization for *E. coli* by Life Technologies [18]. The plasmid pTrc-Sc4CL(M)-VvSTS expresses the synthetic genes described above from the Trc promoter.

### 2.2. Construction of plasmid pTrc-Sc4CL(M)-VvSTS

A 1325-bp DNA fragment containing the gene VvSTS was obtained from the plasmid pTrc-Sc4CL-VvSTS [16] by digestion with the restriction enzymes HindIII/KpnI and gel-purified. The plasmid pTrc-Sc4CL(M)-AhSTS was digested with the restriction enzymes HindIII/KpnI, and the 5694-bp DNA fragment containing the gene Trc promoter, the gene coding for the Sc4CL(M), ampicillin resistance cassette, and plasmid origin of replication was gel-purified and ligated

**Table 1**  
Strains and plasmids used in this study.

	Characteristics	Reference
<i>Strain</i>		
W3110	<i>E. coli</i> F, λ <sup>-</sup> , INV (rrnD <sup>-</sup> rrnE)1	ATCC 27325
W3110Δ-35 <i>fabI</i> ::Cm	W3110 with a deletion of 258 bp upstream of the <i>fabI</i> gene, starting at the -28 position.	This work
<i>Plasmid</i>		
pTrc-Sc4CL(M)-VvSTS	Synthetic codon-optimized genes encoding mutant version of 4CL from <i>S. coelicolor</i> A2 and STS from <i>Vitis vinifera</i> cloned in the expression vector pTrc99A.	[18]

with the 1325-bp fragment. The resultant plasmid pTrc-Sc4CL(M)-VvSTS was verified by restriction and sequence analyses.

### 2.3. Chromosomal deletion of the -35 and upstream promoter regions of the gene *fabI*

The transcriptional start site of the gene *fabI* has been experimentally determined by My et al. [19]. Based on this information, the -10 and -35 promoter sequences were localized. For the deletion of the -35 and upstream regions of the *fabI* promoter, the primers employed were -35*fabI*-F (5'TGCTGGAGAATATTCGACGGAAGTGAACCGCGGTACCCTCTCCCTGATGTGTAGGCTGGAGCTGCTTC3') and -35*fabI*-R (5'GTAAACA GTACGAACAGATAAACGGTTATTATAATCAACCTGGCTGTGAGCATATGAA TATCTCTTAG3'). The 5' end of the primer -35*fabI*-F has 50 bases of homology to the 32' terminal region of the gene *ycjD* up to the stop codon and 20 bases of homology to the P1 sequence of the plasmid pKD3 for amplification of FRT sequence and the chloramphenicol (Cm) resistance cassette [18]. The 5' end of the primer -35*fabI*-R includes 50 bases of homology to the 5' region of the gene *fabI* up to the -28 position of the promoter region and 20 bases of homology to the P2 sequence of the plasmid pKD3. The PCR reaction was performed with standard buffer conditions, Pfu DNA polymerase, one cycle of initial denaturation at 95°C for 3 min, 30 cycles of denaturation at 95°C for 30 s, annealing at 61°C for 60 s, and extension at 68°C for 4 min for a total volume of 50 μL. The PCR product was gel-purified and digested with the restriction enzyme DpnI. The purified PCR DNA from the restriction reaction mixture was purified and electroporated into the strain W3110 that was previously transformed with the plasmid pKD46 [20]. Candidate mutant strains were screened by plating them in Luria-Bertani (LB) solid medium with Cm at a concentration of 30 μg/mL. Colonies displaying Cm resistance were screened by performing PCR with the primers c2-F (5'GATCTTCGTCACAGGTAGG3') [18] and *fabI*11-R (5'TACGGATCGGACCAGCAGAG3') and sequencing the amplified product. A colony displaying the expected chromosomal deletion was named W3110Δ-35*fabI*::Cm.

### 2.4. Culture media and growth conditions

Solid and liquid LB media were used. The antibiotics carbenicillin (Cb; 100 μg/mL) and Cm (30 μg/mL) were used for strain selection. Cultures for pinosylvin production were grown in LB medium supplemented with glycerol 10 g/L. The inoculum for pinosylvin production was started with an overnight 3-mL culture in LB medium supplemented with Cb 100 μg/mL in glass tubes at 37°C and 300 rpm for 12 h. This culture was used to inoculate 50 mL of LB medium supplemented with cinnamic acid 3 mM and Cb 100 μg/mL present in 250-mL baffled shake flasks. The initial O.D.600 of the culture was 0.1, and when it reached an O.D.600 of 0.8, isopropyl-β-D-thiogalactopyranoside was added at a final concentration of 0.5 mM. After 20 h, a sample was taken for pinosylvin quantification. All cultures were performed in triplicate.

## 2.5. RNA extraction, cDNA synthesis, and RT-qPCR analysis

Samples from shaken flask cultures of the strains W3110/pTrc-Sc4CL(M)-VvSTS and W3110Δ-35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS were collected for RNA extraction at the exponential phase. RNA was extracted using hot phenol equilibrated with water, and cDNA synthesis was performed with the RevertAid H First Strand cDNA Synthesis kit (Fermentas, USA) and a mixture of specific DNA primers, as previously reported by Flores et al. [21]. The qPCR experiments were performed with the ABI Prism 7300 Real-Time PCR System (Applied Biosystems, USA) using Maxima SYBRGreen PCR Master Mix (Fermentas, USA) and reaction conditions described previously [21]. The quantification technique used to compare data was the 2- $\Delta\Delta$ CT method [22], and the results were normalized using the *ihfB* gene as an internal control. The qPCR experiments complied with the MIQE guidelines for publication of quantitative real-time PCR experiments [23]. RNA extraction and cDNA synthesis reactions were performed from three independent cultivations for each strain.

## 2.6. Analytical methods

Cell growth was monitored by measuring optical density at 600 nm (O.D.600) in a spectrophotometer (Beckman DU-70; Fullerton, CA, USA). Dry cell weight (DCW) was calculated by multiplying the absorbance at 600 nm with a previously determined coefficient factor of 0.37 g/L [24]. Pinosylvin concentration was determined using an HPLC system (Agilent 1100 System, Agilent Technologies, Palo Alto, CA, USA), with a reverse-phase column (Phenomenex Synergi Hydro RP C18, 150 × 4.6 mm, 4  $\mu$ m, Phenomenex, Torrance, CA, USA) and employing photodiode array detection. Samples were eluted using a gradient in which the mobile phase A was 0.1% TFA in water and the mobile phase B was 0.1% TFA in methanol. Gradient started at 20% B, followed by linear increases to 40% in 5 min, then to 80% in 5 min and then maintained for 3 min. Flow rate was 0.5 mL. Pinosylvin was detected by diode array at  $\lambda = 306$  nm. Pinosylvin (98% purity) used for preparing the standard curve was purchased from Sigma-Aldrich.

## 3. Results and discussion

### 3.1. Construction of an *E. coli* strain with chromosomal deletion of the -35 and upstream promoter regions of the gene *fabI*


A recombinering strategy was employed to delete 258 bp upstream of the *fabI* gene, starting at the -28 position, thus including the -35 promoter sequence [19,20] (Fig. 2). The resulting strain was named W3110Δ-35*fabI*::Cm. To determine whether such modification had an

effect on growth capacity, the strains W3110Δ-35*fabI*::Cm and W3110 were grown using several culture medium compositions. When grown in LB medium, the strains W3110 and W3110Δ-35*fabI*::Cm displayed  $\mu$  values of  $1.32 \pm 0.017$  and  $1.03 \pm 0.010$  h<sup>-1</sup>, respectively. In M9 medium + glucose 10 g/L, the strains W3110 and W3110Δ-35*fabI*::Cm showed  $\mu$  values of  $0.49 \pm 0.001$  and  $0.50 \pm 0.001$  h<sup>-1</sup>, respectively. A 22% reduction in specific growth rate of the mutant strain compared to that of the wild type was observed in LB medium. By contrast, no difference in  $\mu$  values was detected when these two strains were grown in the M9 minimal medium. An explanation for these results is that both strains are considered to grow faster on LB medium; therefore, to sustain such growth rate, a higher cellular demand for fatty acid synthesis is required. Under these conditions, a reduction in the expression of *fabI* gene would cause a reduction in the level of enoyl-acyl carrier protein reductase, thus limiting fatty acid biosynthetic capacity and growth.

### 3.2. Construction and characterization of *E. coli* strains for the production of pinosylvin

To generate *E. coli* strains to produce pinosylvin from cinnamic acid, it is necessary to express genes that encode the 4CL and STS enzymes. In this work, we utilized a synthetic gene that encodes a mutant version of 4-coumaroyl-CoA ligase from *S. coelicolor* A3 (2) having a replacement of amino acid residue alanine at position 294 for a glycine residue (A294G) [25]. This mutant version of 4CL (Ss4CL(M)) displays threefold higher catalytic activity toward cinnamic acid compared to the wild-type enzyme [25]. A synthetic gene encoding STS from *V. vinifera* (VvSTS) was also used in this study because this enzyme has been shown to employ cinnamoyl-CoA as the substrate [12,18]. Genes encoding Ss4CL(M) and VvSTS were cloned as an operon under transcriptional control of the Trc promoter to generate plasmid pTrc-Sc4CL(M)-VvSTS. The strains W3110 and W3110Δ-35*fabI*::Cm were transformed with this plasmid to generate W3110/pTrc-Sc4CL(M)-VvSTS and W3110Δ-35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS. To determine the effect of deleting the -35 and upstream promoter regions of the gene *fabI* on its transcript level in the production strains, RT-qPCR was performed with total mRNA extracted from W3110/pTrc-Sc4CL(M)-VvSTS and W3110Δ-35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS grown in LB + glycerol 10 g/L + 3 mM cinnamic acid. These experiments showed a 60% reduction in *fabI* transcript level in mutant strain W3110Δ-35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS compared to W3110/pTrc-Sc4CL(M)-VvSTS.

Both the -35 and -10 promoter motifs are required to allow recognition by the RNA polymerase carrying sigma 70. However, the -10 sequence is recognized as the most highly conserved and essential




**Fig. 2.** Deletion of *fabI* -35 and upstream region. Chromosomal content of the gene *fabI* in W3110 A) and W3110Δ-35*fabI*::Cm B). The transcription start site for the gene *fabI* is indicated with +1 and the promoter hexamer sequences with -10 and -35.

promoter motif [26,27]. The -10 promoter sequence is particularly important because its recognition drives the initial promoter opening, which is an essential step in transcription initiation [28]. In the gene *fabI*, the -10 promoter motif has the sequence TATAAT, which is identical to the *E. coli* consensus (Fig. 2). By contrast, the -35 motif sequence TAGCTA shows low similarity to the consensus (TTGACA) [27]. Therefore, in this study, the -35 motif of the gene *fabI* was eliminated, expecting to have a negative effect on transcription initiation activity but without the complete elimination of promoter activity.

For the production of pinosylvin, these strains were grown in shake flask cultures in LB medium with glycerol 10 g/L and cinnamic acid

3 mM (0.44 g/L). Under these conditions,  $\mu$  values of  $0.63 \pm 0.001$  and  $0.51 \pm 0.001 \text{ h}^{-1}$  were determined for the strains W3110 and W3110 $\Delta$ -35*fabI*::Cm, respectively. After 23.5 h of culture time, the biomass concentration was  $4.17 \pm 0.09 \text{ g/L}$  for W3110/pTrc-Sc4CL (M)-VvSTS. The strain W3110 $\Delta$ -35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS reached  $3.51 \pm 0.07 \text{ g/L}$  of biomass after 24 h. In these cultures, the final pinosylvin titers corresponded to  $34.89 \pm 4.81$  and  $52.67 \pm 2.95 \text{ mg/L}$  for W3110/pTrc-Sc4CL(M)-VvSTS and W3110 $\Delta$ -35*fabI*::Cm/pTrc-Sc4CL(M)-VvSTS, respectively. Fig. 3 shows an HPLC chromatogram including standards for pinosylvin (panel A) and cinnamic acid (panel B), as well as a sample from a production culture (panel C), where the substrate cinnamic acid and the produced



**Fig. 3.** Chromatograms and associated diode array detection spectra of A: Pinosylvin standard. B: Cinnamic acid standard. C: Production sample at the end point of culture. To confirm peak identity in C, in addition to retention times, diode array detection spectrum was compared to that of the corresponding standards.

pinosylvin were detected and quantified. The calculated yields of pinosylvin produced by unit of biomass ( $Y_{\text{pinosylvin/biomass}}$ ) corresponded to 8.37 and 15 mg pinosylvin/g biomass for W3110/pTrc-Sc4CL(M)-VvSTS and W3110 $\Delta$ -35fabl::Cm/pTrc-Sc4CL(M)-VvSTS, respectively.

#### 4. Conclusion

We report here the construction and characterization of *E. coli* strains for the production of the stilbene pinosylvin from glycerol and cinnamic acid. An approach based on the reduction in the expression of the *fabI* gene was evaluated. A deletion of the *fabI* promoter region resulted in a reduction in transcript level and an increase in pinosylvin production. These results strongly suggest an increased availability of the precursor malonyl-CoA for stilbene biosynthesis in the mutant strain. Based on these results, further efforts toward strain improvement can focus on exploring methods to modulate *fabI* expression by employing synthetic promoters or regulatory circuits with the aim of determining the optimal level of FabI for improving the synthesis of stilbenes and other compounds that use malonyl-CoA as precursor.

#### Ethical approval

This article does not include any studies with human participants or animals performed by any of the authors.

#### Informed consent

This article does not involve any informed consent.

#### Conflict of interest

The authors declare that they have no conflict of interest.

#### Financial support

This work was supported by the Consejo Nacional de Ciencia y Tecnología, México (177,568).

#### Acknowledgments

CS was supported by a fellowship from CONACyT. The authors thank René Ramirez for the generation of pTrc-Ss4CL(M)-AhSTS and also Paul Gaytán, Jorge Yáñez, Eugenio López, and Santiago Becerra for the synthesis of oligonucleotides.

#### References

- Jancinova V, Perecko T, Nosal R, et al. The natural stilbenoid pinosylvin and activated neutrophils: Effects on oxidative burst, protein kinase C, apoptosis and efficiency in adjuvant arthritis. *Acta Pharmacol Sin* 2012;33(10):1285–92. <https://dx.doi.org/10.1038%2Faps.2012.77>.
- Jeandet P, Delaunois B, Conreux A, et al. Biosynthesis, metabolism, molecular engineering and biological functions of stilbene phytoalexins in plants. *Biofactors* 2010;36(5):331–41. <https://doi.org/10.1002/biof.108>.
- Jeandet P, Hébrard C, Deville MA, et al. Deciphering the role of phytoalexins in plant-microorganism interactions and human health. *Molecules* 2014;19(11):18033–56. <https://doi.org/10.3390/molecules191118033>.
- Ribera AE, Zuñiga G. Induced plant secondary metabolites for phytopathogenic fungi control: A review. *J Soil Sci Plant Nutr* 2012;12(4):893–911. <https://doi.org/10.4067/S0718-95162012005000040>.
- Reinisalo M, Kärnlund A, Koskela A, et al. Polyphenol stilbenes: Molecular mechanisms of defence against oxidative stress and aging-related diseases. *Oxid Med Cell Longev* 2015;9:24. <https://doi.org/10.1155/2015/340520>.
- Koskela A, Reinisalo M, Hyttinen JM, et al. Pinosylvin-mediated protection against oxidative stress in human retinal pigment epithelial cells. *Mol Vis* 2014;20:760–9.
- Laavola M, Nieminen R, Leppänen T, et al. Pinosylvin and monomethylpinosylvin, constituents of an extract from the knot of *Pinus sylvestris*, reduce inflammatory gene expression and inflammatory responses in vivo. *J Agric Food Chem* 2015;63(13):3445–53. <https://doi.org/10.1021/jf504606m>.
- Chong JL, Poutaraud A, Huguency P. Metabolism and roles of stilbenes in plants. *Plant Sci* 2009;177(3):143–55. <https://doi.org/10.1016/j.plantsci.2009.05.012>.
- Park SR, Yoon JA, Paik JH, et al. Engineering of plant-specific phenylpropanoids biosynthesis in *Streptomyces venezuelae*. *J Biotechnol* 2009;141(3–4):181–8. <https://doi.org/10.1016/j.jbiotec.2009.03.013>.
- van Summeren-Wesenhagen PV, Marienhagen J. Metabolic engineering of *Escherichia coli* for the synthesis of the plant polyphenol pinosylvin. *Appl Environ Microbiol* 2015;81(3):840–9. <https://doi.org/10.1128/AEM.02966-14>.
- Moche M, Schneider G, Edwards P, et al. Structure of the complex between the antibiotic cerulenin and its target,  $\beta$ -ketoacyl-acyl carrier protein synthase. *J Biol Chem* 1999;274:6031–4. <https://doi.org/10.1074/jbc.274.10.6031>.
- Liang JL, Guo LQ, Lin JF, et al. A novel process for obtaining pinosylvin using combinatorial bioengineering in *Escherichia coli*. *World J Microbiol Biotechnol* 2016;32(6):1–10. <https://doi.org/10.1007/s11274-016-2062-z>.
- Wu J, Du G, Zhou J, et al. Metabolic engineering of *Escherichia coli* for (2S)-pinocembrin production from glucose by a modular metabolic strategy. *Metab Eng* 2013;16:48–55. <https://doi.org/10.1016/j.ymben.2012.11.009>.
- Zha W, Rubin-Pitel SB, Shao Z, et al. Improving cellular malonyl-CoA level in *Escherichia coli* via metabolic engineering. *Metab Eng* 2009;11(3):192–8. <https://doi.org/10.1016/j.ymben.2009.01.005>.
- Heath RJ, Rock CO. Regulation of malonyl-CoA metabolism by acyl-acyl carrier protein and  $\beta$ -ketoacyl-acyl carrier protein synthases in *Escherichia coli*. *J Biol Chem* 1995;270(26):15531–8. <https://doi.org/10.1074/jbc.270.26.15531>.
- Heath RJ, Rock CO. Enoyl-acyl carrier protein reductase (FabI) plays a determinant role in completing cycles of fatty acid elongation in *Escherichia coli*. *J Biol Chem* 1995;270:26538–42. <https://doi.org/10.1074/jbc.270.44.26538>.
- Bachman BJ. Pedigrees of some mutant strains of *Escherichia coli* K12. *Bacteriol Rev* 1972;36(4):525–57.
- Camacho-Zaragoza JM, Hernández-Chávez G, Moreno-Avitia F, et al. Engineering of a microbial coculture of *Escherichia coli* strains for the biosynthesis of resveratrol. *Microb Cell Fact* 2016;15(1):163. <https://doi.org/10.1186/s12934-016-0562-z>.
- My L, Achkar NG, Viala JP, et al. Reassessment of the genetic regulation of fatty acid synthesis in *Escherichia coli*: Global positive control by the dual functional regulator FadR. *J Bacteriol* 2015;197(11):1862–72. <https://doi.org/10.1128/JB.00064-15>.
- Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci* 2000;97(12):6640–5. <https://doi.org/10.1073/pnas.120163297>.
- Flores N, Leal L, Sigala JC, et al. Growth recovery on glucose under aerobic conditions of an *Escherichia coli* strain carrying a phosphoenolpyruvate: Carbohydrate phosphotransferase system deletion by inactivating *arcA* and overexpressing the genes coding for glucokinase and galactose permease. *J Mol Microbiol Biotechnol* 2007;13(1–3):105–16. <https://doi.org/10.1159/000103602>.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* 2001;25(4):402–8. <https://doi.org/10.1006/meth.2001.1262>.
- Bustin SA, Benes V, Garson JA, et al. The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009;55(4):611–22. <https://doi.org/10.1373/clinchem.2008.112797>.
- Lagunas-Muñoz VH, Cabrera-Valladares N, Bolívar F, et al. Optimum melanin production using recombinant *Escherichia coli*. *J Appl Microbiol* 2006;101(5):1002–8. <https://doi.org/10.1111/j.1365-2672.2006.03013.x>.
- Kaneko M, Ohnishi Y, Horinouchi S. Cinnamate: Coenzyme A ligase from the filamentous bacterium *Streptomyces coelicolor* A3 (2). *J Bacteriol* 2003;185(1):20–7. <https://doi.org/10.1128/JB.185.1.20-27.2003>.
- Hook-Barnard IG, Hinton DM. Transcription initiation by mix and match elements: Flexibility for polymerase binding to bacterial promoters. *Gene Regul Syst Biol* 2007;1:275–93. <https://doi.org/10.1177/117762500700100020>.
- Shultzberger RK, Chen Z, Lewis KA, et al. Anatomy of *Escherichia coli*  $\sigma$  70 promoters. *Nucleic Acids Res* 2006;35(3):771–88. <https://dx.doi.org/10.1093%2Fnar%2F35.3.771>.
- Feklistov A. Darst SA structural basis for promoter – 10 element recognition by the bacterial RNA polymerase  $\sigma$  subunit. *Cell* 2011;147(6):1257–69. <https://doi.org/10.1016/j.cell.2011.10.041>.