

Construction and application of a built-in dual luciferase reporter for microRNA functional analysis

Yanzhen Bi¹ ✉ · Xinmin Zheng¹ · Changwei Shao² · Wen Pan³ · Li Jiang² · Huiwu Ouyang²

¹ Hubei Key Laboratory of Animal Embryo Engineering and Molecular Breeding, Institute of Animal Husbandry and Veterinary, Hubei Academy of Agricultural Science, China

² RNA Group, College of Life Science, Wuhan University, China

³ The Third Affiliated Hospital of Sun Yat-Sen University, Tianhe District, China

✉ Corresponding author: biyanzhen@gmail.com

Received October 27, 2010 / Accepted January 19, 2011

Published online: March 15, 2011

© 2011 by Pontificia Universidad Católica de Valparaíso, Chile

ABSTRACT

Background: As key gene regulators, microRNAs post-transcriptionally modulate gene expression via binding to partially complementary sequence in the 3' UTR of target mRNA. An accurate, rapid and quantitative tool for sensing and validation of miRNA targets is of crucial significance to decipher the functional implications of miRNAs in cellular pathways.

Results: Taking advantage of an improved restriction-free cloning method, we engineered a novel built-in dual luciferase reporter plasmid where *Firefly* and *Renilla* luciferase genes were assembled in a single plasmid named "pFila". This design eliminates the transfection of a separate control plasmid and thus minimizes the time and labor required for miRNA-target sensing assays. pFila consistently produces *Firefly* and *Renilla* luciferase activities when transfected into human-, monkey- and mouse-derived mammalian cell systems. Moreover, pFila is capable of recapitulating the interaction of miR-16 and its known target CCNE1 in Hela cells. Additionally, pFila is shown to be a sensitive miR-biosensor by evaluating the inhibition efficiency of endogenous miRNA.

Conclusions: pFila would facilitate miRNA target identification and verification in a rapid and simplified manner. Also, pFila is a sensitive biosensor for active miRNA profiling *in vivo*.

Keywords: biosensor, luciferase, ligase-independent, miRNA, target

INTRODUCTION

MicroRNAs are key post-transcriptional regulators of gene expression in a variety of cellular events. They mediate translational repression, and sometimes destabilization, of target mRNAs by directing miRISC (microRNA-induced silencing complex) to imperfect complementary sequences in 3' UTR (Bartel, 2009). It has been predicted that more than 60% of human genes are putative targets of one or more miRNAs, while it is also suggested that an individual miRNA is capable of regulating multiple target mRNAs (Backes et al. 2010). Consequently, a major challenge in miRNA study is the experimental identification and validation of its functional target(s). Various reporter systems have been developed to probe the interaction between individual miRNA and its target (Lee et al. 2008), of which dual luciferase assay is widely adopted to achieve this end (Brennecke et al. 2005). The current dual luciferase assay encompasses two separate plasmids, one containing the region of interest and the other serving as internal control to normalize transfection variation (Robertson et al. 2010). A major drawback of this system is the tedious steps for preparing and transfecting control vector. A more convenient reporter is desired to simplify the conventional protocol.

In addition, there are several methods, including northern blotting, real-time PCR, microarray and deep sequencing, to quantify miRNAs (Willenbrock et al. 2009). However, these techniques simply output the homeostasis of endogenous miRNAs other than active molecules. Moreover, these approaches are either time-consuming and laborious or cost-ineffective. Therefore, a convenient and quantitative miRNA biosensor is desired to measure functional miRNA *in vivo*.

In this study, by taking advantage of ligase-free homologous recombination in *E. coli*, we engineered a novel reporter that integrated *Firefly* and *Renilla* luciferase genes (*Fluc* and *Rluc*, same below) in a single plasmid. Its expressivity and applicability were further examined to demonstrate that this novel reporter will facilitate the screening and sensing of miRNAs and their targets in a simplified and precise manner.

MATERIALS AND METHODS

DNA and RNA oligos

Unless stated elsewhere, all DNA and RNA oligos are presented as 5'→3' direction. Primers for amplifying *Fluc* gene were Pf, **AAGGATCCAGGTGGCACTTTTCG TGCGATCTGCATCTCAATTAG**; Pr, GAAAAATAAACAAATAGGGGTTCCGCGCAC **CTCACATGTTCTTTCCTGC** (sequence annealing to *Fluc* gene was shown in boldface; sequence complementary to insertion site on pRL-TK was underlined). 3' outermost primer P2R was CGAAAAGTGCCACCTGGATCCTT. Sequencing primers for pFila was SF, GATGCACCTG ATGAAATGGG; SR, AGGACAGGTG CCGGCAGCGC. For creation of *Apal* site, see details in reference (Wang et al. 2009). RNA oligos were chemically synthesized and purified by Genepharma Co. Ltd., (Shanghai, P.R. China). Human miR16-1 was sense UAGCAGCAGUAAAUUUGGCG and antisense CGCCAAUUAUUACGUGCUGCUA. Negative control for miRNA mimics was sense UUGUACUACACAAAAGUACUG and antisense CAGUACUUUUGUGUAGUACAA. siRNA against *Rluc* mRNA was sense GUAGCGGGUGUAUUUACdTdT and antisense GUAUAAUACCGCGCUACdTdT. Methylated anti-miR-16-1 inhibitor CGC CAA UAU UUACGU GCU GCU A, scramble anti-miR control UUG UAC UAC ACA AAA GUA CUG.

Construction of plasmids

pFila was fabricated by bridging-PCR coupled with homologous recombination in bacteria. A duplex bridging PCR was conducted in a 50 µl mixture: Pf 4 µl (250 nM), Pr 2 µl (125 nM), P2R primer 2 µl (125 nM), pGL3-promoter plasmid 1 µl (5 ng), modified pRL-TK plasmid 1 µl (10 ng or 50 ng), 2 mM dNTP 5 µl, 25 mM MgSO₄ 2 µl, 10 x KOD buffer 5 µl, KOD plus 1 µl (1 unit), PCR-grade water 27 µl. The condition was: 95°C 2 min, 30 cycles of (95°C 15 sec, 55°C 30 sec, 68°C 6.5 min). The PCR products were digested with *DpnI* (Fermentas, Lithuania) at 37°C for 2 hrs to destroy methylated plasmids while keeping the nascent DNA intact with the following reaction: PCR products, 26 µl; 10 x Tango buffer 3 µl; *Dpn I* 1 µl (1 unit). An aliquot of 5 µl digested PCR products were transformed into *E. coli* DH5α to generate recombinants that were subsequently sequenced to verify the integrity. Pf and Pr primer pair was used to amplify *Fluc* gene composed of SV40 promoter, *Fluc* coding region and SV40 late poly(A) signal. *Fluc* gene was designed to fuse into a modified pRL-TK plasmid downstream of 3'UTR of *Rluc* and upstream of beta-lactamase gene. For the sequence context of human CCNE1 3'UTR, see details in reference (Wang et al. 2009). The wild-type and mutated human CCNE1 target regions were sub-cloned into pFila with *Xba I* and *Apal I*.

Cell culture, transfection and dual luciferase assay

Human cervical carcinoma Hela cells, African green monkey kidney Vero cells and mouse myoblast C2C12 cells were maintained in high glucose DMEM (Invitrogen) supplemented with 10% fetal calf serum (Gibco) at 37°C and 5% CO₂. 4 × 10⁴ cells were seeded in a 24-well plate one day before transfection. For miRNA mimics and plasmid co-transfection, 1 µl 20 µM chemically synthesized miR16 mimics and 50 ng pFila (pFila-CCNE1-wildtype and pFila-CCNE1-mut1&2) or 50 ng pGL3-promoter (internal control) and pRL-ML plasmids (pRL-ML-CCNE1-wildtype and pRL-ML-CCNE1-mut1&2; 25 ng each) were mixed with 2 µl Lipofectamine2000 (Invitrogen) as transfection complex. For evaluation of endogenous miRNA inhibition with pFila, 20 nM 2'-O-methylated anti-human miR16-1 inhibitor was transfected into Hela cells by 0.5 µl Lipofectamine2000 (Invitrogen), RNAiMAX (Invitrogen), Sofast

(Sunmabio, China), Fugene (Roche), respectively. All transfections were performed in three independent experiments with each in triplicate. A DLRTM Assay (Promega) was adopted to measure luciferase activity in a Glomax luminometer essentially according to manufacturer's instruction.

Statistical analysis

Luciferase levels were reported as ratio over that observed in control transfections, where *Rluc* activities were normalized to *Fluc* activities. The data represented the mean \pm S.D. of three independent experiments and were analyzed by Student's *t*-test. Differences below $p < 0.01$ were regarded as significant.

Accession number

The sequence and annotation of pFila has been deposited in Genbank with accession number HQ425563. pFila is freely available upon request.




Fig. 1 Concept and engineering of pFila. (a) In this reaction, the primary process was the production of *Fluc* gene by Pf (Pf is composed of P1 and P2, which are complementary to upstream of *Fluc* gene and pRL-TK plasmid, respectively) and Pr (Pr is composed of P3 and P4, which are complementary to downstream of *Fluc* gene and pRL-TK plasmid, respectively) primers. In the secondary process, amplified *Fluc* gene bearing a deliberately designed region annealed with its homologous sequence in pRL-TK, thus generating a large linear fragment. In order to exponentially amplify the linear fragment, the outermost Pf and P2R (complementary to P2) primer pair initiated the ternary process to produce adequate fused fragment. (b) Duplex bridge PCR. The 2.4 kb fragment was *Fluc* gene, while the 6.4 kb fragment was the desired fusion product. (c) Map of pFila. The engineered pFila is 6486 bp in length. *Xba*I and *Apa*I restriction sites are located in the 3'UTR of *Rluc* to facilitate the cloning of miRNA target for reporter assay. (d) 5' and 3' seaming sites of *Fluc* into pRL-TK. For detailed information, please refer to Gene sequence 1.

RESULTS AND DISCUSSION

The complicated procedures of the current dual luciferase reporter assay for miRNA target screening prompted us to upgrade its practicality for simplified manipulations. Specifically, we aimed to integrate *Fluc* and *Rluc* genes in a single vector. As restriction sites were not available to sub-clone *Fluc* gene to pRL-TK plasmid with ligase-dependent method, we adopted an improved restriction-free gene fusion approach inspired by the principle of site-directed mutagenesis (Zheng et al. 2004). As shown in Figure 1a and Figure 1b, a duplex bridging PCR was carried out to produce the 6.4 kb linear fusion fragment

with homologous sequences at both 5' and 3' ends. The PCR products were digested by *DpnI* and transformed into *E. coli* to achieve the circular plasmid based on homologous recombination. Sequencing of the recombinant (named pFila) revealed that *Fluc* gene had been successfully fused into pRL-TK plasmid at designed location (Figure 1c and Figure 1d; Gene sequence 1).

We then evaluated the expressivity of pFila in different mammalian systems. As presented in Figure 2a, luminescence of pFila was reported in a wide linear range when transfected into human-, mouse- and monkey-sourced cell lines at gradient amounts, indicating that pFila consistently produces luciferases *in vivo*. This also implies that the ordered assembly of *Fluc* and *Rluc* luciferase genes in pRL-TK plasmid does not interfere with their individual expression. Next, the applicability and reproducibility of pFila were examined by recapitulating the regulation of human miR16-1 and its known target CCNE1 (Wang et al. 2009). miR16-1 mimics down-regulated the *Rluc* activity fused with wild-type CCNE1 3'UTR but not a mutant 3'UTR (Figure 2b); the latter carried altered residues that were introduced in the miR16-1 "seed-pairing" recognition site (Figure 2b). This observation perfectly photocopied the result that was achieved by traditional dual reporter assay (Figure 2b), indicating that pFila as a more convenient reporter is fully applicable to miRNA functional analysis. Finally, we applied pFila carrying the wild-type and mutant 3'UTR of CCNE1 to assessing the blockage efficiency of endogenous miR16-1. 2'-O-methylated anti-miRNA-16-1 RNA oligo was transfected by four types of transfection reagents, *i.e.* Lipofectamine2000, RNAiMAX, Fugene, Sofast. Inhibition efficacy of endogenous miR16-1 varied, of which RNAiMAX achieved the most potent blocking effect. This assay implies that pFila is a sensitive miRNA biosensor to reflect the level of functional miRNAs. It also suggests that the choice of delivery method is an important determinant when conducting loss-of-function analysis of miRNAs.

In summary, we have successfully engineered a novel dual luciferase plasmid that incorporated *Fluc* and *Rluc* genes in a single vector, allowing the simultaneous expression of both luciferase genes. This improvement maintains comparable reproducibility but minimizes the time and labor required in conventional dual luciferase protocol. Furthermore, several lines of evidence were presented to demonstrate its application in miRNA functional analysis. These results indicate that pFila will find its wide application in the screening, identification and validation of miRNA with its potential mRNA targets.




Fig. 2 Expressivity and application of pFila. (a) pFila consistently produces *Fluc* and *Rluc* activities in human-, mouse- and monkey-sourced mammalian cell lines. pFila was gradually transfected into the cells stated above and *Fluc* and *Rluc* luciferase levels were determined by a DLR™ Assay (Promega). As shown in this graph, *Fluc* and *Rluc* activities were in a linear range for all the selected points. (b) pFila is capable of recapitulating the interaction of miR16 and its known target CCNE1. Wild-type and mutant CCNE1 3'UTRs were sub-cloned into pFila and co-transfected with miR16-1 mimics. Conventional dual luciferase reporter was conducted in parallel to compare their reproducibility. siRNA against *Rluc* serves as positive control. *Rluc* with mutant CCNE1 3'UTR was rescued in comparison with pFila-CCNE1-3'UTR-wildtype, indicating that CCNE1 is a direct target of miR16 as previously reported (Wang et al. 2009). (c) miR16-1 inhibitors were co-transfected with pFila-CCNE1-3'UTR-wildtype or mutant plasmids into Hela cells by different transfection reagents to block endogenous miR16-1. Inhibition efficacy varies with the selected agents, where RNAiMAX achieved the most potent inhibition. It implies that pFila is a sensitive biosensor for functional miRNA profiling.

Gene sequence 1. Sequencing of the seaming sites: sequence annealing to Fluc gene was shown in boldface; sequence complementary to insertion site on pRL-TK was underlined.

Seq.1 (sequencing primer SF: GATGCACCTGATGAAATGGG)

TTTTTAAATCCAATTCGGTTGAACGAGTTCTCAAAAATGAACAATAATTCTAGATTCCGAGATATCGGTAATGGGC
 CCTAGAGCGCCGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGACAAACCACAACCTAGAATGCAGTGA
 AAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACCAAGTTA
 ACAACAACAATTGCATTCAATTTATGTTTCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAAACCT
 CTACAAATGTGGTAAATTCGATAAAGGATCCAGGTGGCACTTTTCG**TGCGATCTGCATCTCAATTAGT**CAGCAACCA
 TAGTCCCGCCCTAACTCCGCCCATCCCGCCCTAACTCCGCCAGTTCCGCCCATTTCTCCGCCCATCGCTGACT
 AATTTTTTTTTATTTATGACAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTT
 GGAGGCTTAGGCTTTTGCAAAAAGCTTGGCATTCCGGTACTGTTGGTAAAGCCACCATGGAAGACGCCAAAACAT
 AAAGAAAGGCCCGGCCCATTTCTATCCGCTGGAAGATGGAACCGCTGGAGAGCAACTGCATAAGGCTATGAAGAGA
 TACGCCCTGGTTCCGGAAACAATTGCTTTTACAGATGCACATATCGAGGTGGACATCACTTACGCTGAGTACTTCG
 AAATGTCCGTTCCGGTTGGCAGAAGCTATGAAACGATATGGGCTGAATACAAATCACAGAATCGTCGTATGCAGTGA
 AAATCTCTTCAATTCTTTATGCCGCTGTTGGGCGGTTATTTATCGGAGTTGCAGTTGCCCGCGAACGACAT
 TATAATGAACGTAATTGCTCAACAGTATGGGCATTTCCGAGCCTACCGTGGTGTTCGTTTCCAAAAGGGTTGCA
 AAAAAATTTGAAGCTGCAAAAAGCTCCCAATCATCCAAAATTTATTTATCATGGATTCTAAAACGGATTACCAGG
 GGATTCAGTCGATGTACACGTCGTCACATCTCATCTACCTTCCCGTTTTATGATACGATTTTGTGCAAGAGCTTTC
 GATAGACAAGACAATTGCACTGATCATGACCTCTCTGATCTACTGGACTGCTAGGTGCGACCTTGTCTCATAGAC
 TGCTTGCCGGTAGAAATTCCTCGCC

Seq.2 (sequencing primer SR: AGGACAGGTG CCGGCAGCGC)

CCTGGAAATAAGAGGTGAGGTGATCATCAGGTAATGGGGTACTGTGACGATACCCAATGCTATCAGGAGGCACGT
 TATTACGCGTCTGTCTATTCGGTCATCATAGTTGCTGACTCCCGTCGGTAGATAACTACGATACGGAGGCTACAA
 TCTGCCAGTGTGCATGATACGCGAGACCAGCTCACCGGCTCAGATTATCAGCATAACAGCAGCGAGGGCCGAG
 CGCAGAGTGTCTGCATTTATCCGCTCCATCCAGTCTATAATGTGCGGGAAAGCTAGAGTAAGTAGTTCGCCAGTTAA
 TAGTTTGGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCGTTCGTTGGTATGGCTTCATTACG
 TCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCCTC
 CGATCGTTGTGAGAAGTAAGTTGGCCGAGTGTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGT* /
 CATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGTACTCAACCAAGTCATTCTGAGAAATAGTGTATCGCGCGA
 CCGAGTTGCTCTTCCCGCGCGTCAATACGGGATAATACCGGCCACATAGCAGAACTTTAAAAGTGTCTATCATTG
 GAAAACGTTCTTCGGGCGCAAAACTCTCAAGGATCTTACCCTGTGAGATCCAGTTCGATGTAACCCACTCGTGC
 ACCCAACTGATCTTACGATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCA
 AAAAAAGGAAATAAGGCGCACACGGAAATGTTGAATACTATACTCTTCTTTTCAATATATTGAAGCATTTATC
 AGGGTTATTGCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACCTC
ACATGTTCTTTCCTGCGTTATCCCTGATTTCTGTGGATAAACCCTATTACCCTTTGAGTGAGCTGATACCGCTCG
 CCGCAGCCGAACGACCGAGCGCAGCGAGTCAAGCGAGTGGG

Seq.3 pFla sequence

Base pair:	6486bp
HSV TK promoter	7-759
Chimeric intron	826-962
T7 RNA polymerase Promoter (-17 to +2)	1006-1024
T7 RNA polymerase transcription initiation site	1023
Rluc reporter gene	1034-1969
XbaI restriction site	1971-1976
Apal restriction site	1995-2000
SV40 late polyadenylation signal(upstream)	2039-2240
SV40 promoter	2279-2481
Fluc reporter gene	2511-4163
SV40 late polyadenylation signal(downstream)	4195-4416
Beta-lactamase (AmpR)	4800-5660
pBR322 plasmid replication origin	5802-6445

1 AGATCTAAAT GAGTCTTCGG ACCTCGCGGG GGCCGCTTAA GCGGTGGTTA
 51 GGGTTTGTCT GACGCGGGGG GAGGGGGAAG GAACGAAACA CTCTCATTCTG
 101 GAGGCGGCTC GGGGTTTGGT CTTGGTGGCC ACGGGCACGC AGAAGAGCGC
 151 CGCGATCCTC TTAAGCACCC CCCC GCCCTC CGTGGAGGCG GGGGTTTGGT
 201 CGGCGGGTGG TAACTGGCGG GCCGCTGACT CGGGCGGGTC GCGCGCCCCA
 251 GAGTGTGACC TTTTCGGTCT GCTCGCAGAC CCCC GGCGCG CGCCGCGCG
 301 GCGGCGACGG GCTCGCTGGG TCCTAGGCTC CATGGGACG GTATACGTGG
 351 ACAGGCTCTG GAGCATCCGC ACGACTGCGG TGATATTACC GGAGACCTTC
 401 TGCGGGACGA GCCGGTCTAC GCGGCTGACG CGGAGCGTCC GTTGGGCGAC

451 AAACACCAGG ACGGGGCACA GGTACTACTAT CTTGTACACC GGAGGCGCGA
501 GGGACTGCAG GAGCTTCAGG GAGTGGCGCA GCTGCTTCAT CCCCCTGGCC
551 CGTTGCTCGC GTTTGCTGGC GGTGTCCCCG GAAGAAATAT ATTTGCATGT
601 CTTTAGTTCT ATGATGACAC AAACCCCGCC CAGCGTCTTG TCATTGGCGA
651 ATTCGAACAC GCAGATGCAG TCGGGGCGGC GCGGTCCCAG GTCCACTTCG
701 CATATTAAGG TGACGCGTGT GGCCTCGAAC ACCGAGCGAC CCTGCAGCGA
751 CCCGCTTAAA AGCTTGATTG TTCTGACACA ACAGTCTCGA ACTTAAGCTG
801 CAGAAGTTGG TCGTGAGGCA CTGGGCAGGT AAGTATCAAG GTTACAAGAC
851 AGGTTTAAAG AGACCAATAG AAACGGGCT TGTCGAGACA GAGAAGACTC
901 TTGCGTTTCT GATAGGCACC TATTGGTCTT ACTGACATCC ACTTTGCCTT
951 TCTCTCCACA GGTGTCCACT CCCAGTTCAA TTACAGCTCT TAAGGCTAGA
1001 GTACTTAATA CGACTCACTA TAGGCTAGCC ACCATGACTT CGAAAGTTA
1051 TGATCCAGAA CAAAGGAAAC GGATGATAAC TGGTCCGCG TGGTGGGCCA
1101 GATGTAACA AATGAATGTT CTTGATTTCAT TTATTAATA TTATGATCA
1151 GAAAAACATG CAGAAAAATG TGTTATTTTT TTACATGGTA ACGCGGCCTC
1201 TCTTTATTTA TTGGCGACATG TTGTGCCACA TATGAGCCA GTAGCGCGGT
1251 GTATTATACC AGACCTTATT GGTATGGGCA AATCAGGCAA ATCTGGTAAT
1301 GGTTCCTATA GGTTACTTGA TCATTACAAA TATCTTACTG CATGGTTTGA
1351 ACTTCTTAAT TTACCAAGA AGATCATTTT TGTCGGCCAT GATTGGGGTG
1401 CTGTTTGGC ATTTTATTAT AGCTATGAGC ATCAAGATA GATCAAAGCA
1451 ATAGTTCACG CTGAAAGTGT AGTAGATGTG ATTGAATCAT GGGATGAATG
1501 GCGTGAATTT GAAGAAGATA TTGCGTTGAT CAAATCTGAA GAAGGAGAAA
1551 AAATGGTTTT GGAGAATAAC TTCTTCGTGG AAACCATGTT GCCATCAAAA
1601 CTCATGAGAA AGTTAGAACC AGAAGAATTT GCAGCATATC TTGAACCATT
1651 CAAAGAGAAA GGTGAAGTTC GTCGTCCAAC ATTATCATGG CCTCGTGAAA
1701 TCCCGTTAGT AAAAGGTGGT AAACCTGACG TTGTACAAAT TGTTAGGAAT
1751 TATAATGCTT ATCTACGTGC AAGTATGATG TTACCAAAAA TGTTTATTGA
1801 ATCGGACCCA GGATTTCTTTT CCAATGCTAT TGTGGAAGGT GCCAAGAAGT
1851 TTCCTAATAC TGAATTTGTC AAAGTAAAAG GTCTTCAATTT TCGCAAGAAA
1901 GATGCACCTG ATGAAATGGG AAAATATATC AAATCGTTTCG TTGAGCGAGT
1951 TCTCAAAAAA GAACAATAAT TCTAGATTCC GAGATATCGG TAATGGGCCC
2001 TAGAGCGGCC GCTTCGAGCA GACATGATAA GATACATTGA TGAGTTTGGG
2051 CAAACCACAA CTAGAATGCA GTGAAAAAAA TGCTTTATTT GTGAAATTTG
2101 TGATGCTATT GCTTTATTTG TAACCATTAT AAGCTGCAAT AAACAAGTTA
2151 ACAACAACAA TTGCATTTCAT TTTATGTTTC AGGTTTCAGG GGAGGTGTGG
2201 GAGGTTTTTT AAAGCAAGTA AAACCTCTAC AAATGTGGTA AAATCGATAA
2251 GGATCCAGGT GGCACTTTTT GTGCGATCTG CATCTCAATT AGTCAGCAAC
2301 CATAGTCCCG CCCCTAACTC CGCCCATCCC GCCCTAACT CGGCCAGTT
2351 CCGCCCATTC TCCGCCCATC CGCTGACTAA TTTTTTTTTT TTATGCAGAG
2401 CCGGAGGCC CCTCGGCCTC TGAGCTATTG CAGAAGTAGT GAGGAGGCTT
2451 TTTTGGAGGC CTAGGCTTTT GCAAAAAGCT TGGCATTCCG GTACTGTTGG
2501 TAAAGCCACC ATGGAAGACG CCAAAAACAT AAAGAAAGGC CCGGCGCCAT
2551 TCTATCCGCT GGAAGATGGA ACCGCTGGAG AGCAACTGCA TAAGGCTATG
2601 AAGAGATACG CCTTGGTTCC TGGAAACAAT GCTTTTACAG ATGCACATAT
2651 CGAGGTGGAC ATCACTTACG CTGAGTACTT CGAAATGTCC GTTCGGTTGG
2701 CAGAAGCTAT GAAACGATAT GGGCTGAATA CAAATCACAG AATCGTCGTA
2751 TGCAGTGAAT ACTCTCTTCA ATTTCTTTATG CCGGTGTTGG GCGGTTTATT
2801 TATCGGAGTT GCAGTTGCGC CCGGAACGCA CATTTATAAT GAACGTGAAT
2851 TGCTCAACAG TATGGGCATT TCGCAGCCTA CCGTGGTGTG GCTTTCCAAA
2901 AAGGGGTTGC AAAAAATTTT GAACGTGCAA AAAAGCTCC CAATCATCCA
2951 AAAAAATATT ATCATGGATT CTAAAACGGA TTACCAGGGA TTTCAGTCGA
3001 TGTACACGTT CGTCACATCT CATCTACCTC CCGTTTTTAA TGAATACGAT
3051 TTTGTGCCAG AGTCCTTCGA TAGGGACAAG ACAATTGCAC TGATCATGAA
3101 CTCTCTGGA TCTACTGGTC TGCCATAAAG TGTCGCTCTG CCTCATAGAA
3151 CTGCCTGCGT GAGATTCTCG CATGCCAGAG ATCCTATTTT TGGAATCAA
3201 ATCATTCGGG ATACTGCGAT TTTAAGTGTG GTTCCATTCC ATCACGGTTT
3251 TGGAAATGTTT ACTACACTCG GATATTTGAT ATGTGGATTG CGAGTCGTCT
3301 TAATGTATAG ATTTGAAGAA GAGCTGTTTC TGAGGAGCCT TCAGGATTAC
3351 AAGATTCAAA GTGCGCTGCT GGTGCCAACC CTATTTCTCT TCTTCGCCAA
3401 AAGCACTCTG ATTGACAAAT ACGATTTATC TAATTTACAC GAAATGCTT
3451 CTGGTGGCGC TATGGCTCTT AAGGAAGTCC GGGGAAGCGG TGCCAAGAGG
3501 TTCCATCTGC CAGGTATCAG GCAAGGATAT GGGCTCACTG AGACTACATC
3551 AGCTATCTG ATTACACCCG AGGGGGATGA TAAACCGGCC GCGGTCCGTA
3601 AAGTTGTTCC ATTTTTTGAA GCGAAGGTTG TGGATCTGGA TACCGGGAAA
3651 ACGCTGGGCG TTAATCAAAG AGGCGAAGTCT TGTGTGAGAG TGTCTATGAT
3701 TATGTCGGGT TATGTAAACA ATCCGGAAGC GACCAACGCC TTGATTGACA
3751 AGGATGGATG GCTACATTCT GGAGACATAG CTTACTGGGA CGAAGACGAA
3801 CACTTCTTCA TCGTTGACCG CCTGAAGTCT CTGATTAAGT ACAAAGGCTA
3851 TCAGGTGGCT CCGCTGAAT TGGAATCCAT CTTGCTCCA CACCCCAACA
3901 TCTTCGACGC AGGTGTCGCA GGTCTTCCCG ACGATGACGC CGGTGAACCT
3951 CCCGCCCGG TTGTTGTTTT GGAGCACGGA AAGACGATGA CGAAAAAGA

4001 GATCGTGGAT TACGTCGCCA GTCAAGTAAC AACCGCGAAA AAGTTGCGCG
 4051 GAGGAGTTGT GTTTGTGGAC GAAGTACCGA AAGGTCTTAC CGGAAAACCTC
 4101 GACGCAAGAA AAATCAGAGA GATCCTCATA AAGGCCAAGA AGGGCGGAAA
 4151 GATCGCCGTG TAATTCTTGA GTCGGGGCGG CCGGCCGCTT CGAGCAGACA
 4201 TGATAAGATA CATTGATGAG TTTGGACAAA CCACAACCTAG AATGCAGTGA
 4251 AAAAAATGCT TTATTTGTGA AATTTGTGAT GCTATTGCTT TATTTGTAAC
 4301 CATTATAAGC TGCAATAAAC AAGTTAACAA CAACAATTGC ATTCATTTTA
 4351 TGTTTCAGGT TCAGGGGGAG GTGTGGGAGG TTTTTTAAAG CAAGTAAAAA
 4401 CTCTACAAAT GTGGTAAAAT CGATAAGGAT CCGTCGACCG ATGCCCTTGA
 4451 GAGCCTTCAA CCCAGTCAGC TCCTTCCGGT GGGCGCGGGG CATGACTATC
 4501 GTCGCCGCAC TTATGACTGT CTTCTTTATC ATGCAACTCG TAGGACAGGT
 4551 GCCGGCAGCG CCTTCCGCT TCCTCGTCA CTGACTCGCT GCGCTCGGTC
 4601 GTTCGGCTGC GGCAGCGGT ATCAGCTCAC TCAAAGGCGG TAATACGGTT
 4651 ATCCACAGAA TCAGGGGATA ACGCAGGAAA GAACATGTGA GGTGCGCGGA
 4701 ACCCCTATTT GTTTATTTT CTAAATACAT TCAAATATGT ATCCGCTCAT
 4751 GAGACAATTAT CCCTGATAAA TGCTTCAATA ATATTGAAAA AGGAAGAGTA
 4801 TGAGTATTCA ACATTTCCGT GTCGCCCTTA TTCCCTTTTT TGCGGCATTT
 4851 TGCCTTCCCTG TTTTGTGCTA CCCAGAAACG CTGGTGAAAG TAAAAGATGC
 4901 TGAAGATCAG TTGGGTGCAC GAGTGGGTTA CATCGAAGT GATCTCAACA
 4951 GCCGTAAGAT CCTTGAGAGT TTTGCGCCCG AAGAACGTTT TCCAATGATG
 5001 AGCACTTTTA AAGTTCTGCT ATGTGGCGCG GTATTATCCC GTATTGACGC
 5051 CGGGCAAGAG CAACTCGGTC GCCGCATACA CTATTCTCAG AATGACTTGG
 5101 TTGAGTACTC ACCAGTCACA GAAAAGCATC TTACGGATGG CATGACAGTA
 5151 AGAGAATTAT CCAGTGCATG CATAACCATG AGTGATAACA CTGCGGCCAA
 5201 CTTACTTCTG ACAACGATCG GAGGACCGAA GGAGCTAACC GCTTTTTTGC
 5251 ACAACATGGG GGATCATGTA ACTCGCCTTG ATCGTTGGGA ACCGGAGCTG
 5301 AATGAAGCCA TACCAAACGA CGAGCGTAC ACCACGATGC CTGTAGCAAT
 5351 GGCACAACG TTGCGCAAAC TATTAAGTGG CGAAGTACT ACTCTAGTCT
 5401 CCCGGCAACA ATTAATAGAC TGGATGGAGG CGGATAAAGT TGCAGGACCA
 5451 CTTCTGCGCT CGGCCCTTCC GGCTGGCTGG TTTATTGCTG ATAAATCTGG
 5501 AGCCGGTGAG CGTGGGTCTC GCGGTATCAT TGCAGCACTG GGGCCAGATG
 5551 GTAAGCCCTC CCGTATCGTA GTTATCTACA CGACGGGGAG TCAGGCAACT
 5601 ATGGATGAAC GAAATAGACA GATCGCTGAG ATAGGTGCCT CACTGATTAA
 5651 GCATTGGTAA CTGTCAGACC AAGTTTACTC ATATATACTT TAGATTGATT
 5701 TAAAACCTCA TTTTAAATTT AAAAGGATCT AGGTGAAGAT CCTTTTTGAT
 5751 AATCTCATGA CCAAATCCC TTAACGTGAG TTTTCGTTCC ACTGAGCGTC
 5801 AGACCCCGTA GAAAAGATCA AAGGATCTT TTGAGATCCT TTTTTTCTGC
 5851 GCGTAATCTG CTGCTTGCAA AAAAAAAC CACCGCTACC AGCGGTGGTT
 5901 TGTTTGCCGG ATCAAGAGCT ACCAAGTCTT TTTCCGAAGG TAACTGGCTT
 5951 GAGCAGAGCG CAGATACCAA ATACTGTTCT TCTAGTGTAG CCGTAGTTAG
 6001 GCCACCATT CAAGAAGTCT GTAGCACC GCCTACATACCT CGCTCTGCTA
 6051 ATCTGTTTAC CAGTGGCTGC TGCCAGTGGC GATAAGTTCGT GTCCTTACCGG
 6101 GTTGACTCA AGACGATAGT TACCGGATAA GGCGCAGCGG TCGGGCTGAA
 6151 CGGGGGTTC GTGCACACAG CCCAGCTTGG AGCGAACGAC CTACACCGAA
 6201 CTGAGATACC TACAGCGTGA GCTATGAGAA AGCGCCACGC TTCCCGAAGG
 6251 GAGAAAGCGG GACAGGTATC CGGTAAGCGG CAGGGTCGGA ACAGGAGAGC
 6301 GCACGAGGGA GCTTCCAGGG GGAACGCTT GGTATCTTTA TAGTCTGTCT
 6351 GGGTTTCGCC ACCTCTGACT TGAGCGTCA TTTTGTGAT GCTCGTCAGG
 6401 GGGGCGGAGC CTATGGAAAA ACGCCAGCAA CGCGGCTTTT TTACGGTTCC
 6451 TGGCCTTTTG CTGCCTTTT GCTCACATGG CTCGAC

ACKNOWLEDGMENTS

We appreciated the technical assistance from Cell Signaling Lab of Wuhan University.

Financial support: This work was funded by a research grant from Hubei Key Laboratory of Animal Embryo Engineering and Molecular Breeding to Y. Z. Bi (2010ZD163).

REFERENCES

- BACKES, C.; MEESE, E.; LENHOF, H.P. and KELLER, A. (2010). A dictionary on microRNAs and their putative target pathways. *Nucleic Acids Research*, vol. 38, no. 13, p. 4476-4486. [\[CrossRef\]](#)
- BARTEL, D.P. (2009). MicroRNAs: Target recognition and regulatory functions. *Cell*, vol. 136, no. 2, p. 215-233. [\[CrossRef\]](#)

- BRENNECKE, J.; STARK, A.; RUSSELL, R.B. and COHEN, S.M. (2005). Principles of microRNA-target recognition. *PLoS Biology*, vol. 3, no. 3, p. e85. [\[CrossRef\]](#)
- LEE, J.Y.; KIM, S.; HWANG, D.W.; JEONG, J.M.; CHUNG, J.K.; LEE, M.C. and LEE, D.S. (2008). Development of a dual-luciferase reporter system for *in vivo* visualization of microRNA biogenesis and posttranscriptional regulation. *The Journal of Nuclear Medicine*, vol. 49, no. 2, p. 285-294. [\[CrossRef\]](#)
- ROBERTSON, B.; DALBY, A.B.; KARPILOW, J.; KHVOROVA, A.; LEAKE, D. and VERMEULEN, A. (2010). Specificity and functionality of microRNA inhibitors. *Silence*, vol. 1, no. 1, p. 10. [\[CrossRef\]](#)
- WANG, F.; FU, X.D.; ZHOU, Y. and ZHANG, Y. (2009). Down-regulation of the cyclin E1 oncogene expression by microRNA-16-1 induces cell cycle arrest in human cancer cells. *BMB Report*, vol. 42, no. 11, p. 725-730.
- WILLENBROCK, H.; SALOMON, J.; SØKILDE, R.; BARKEN, K.B.; HANSEN, T.N.; NIELSEN, F.C.; MØLLER, S. and LITMAN, T. (2009). Quantitative miRNA expression analysis: Comparing microarrays with next-generation sequencing. *RNA*, vol. 15, no. 11, p. 2028-2034. [\[CrossRef\]](#)
- ZHENG, L.; BAUMANN, U. and REYMOND, J.L. (2004). An efficient one-step site-directed and site-saturation mutagenesis protocol. *Nucleic Acids Research*, vol. 32, no. 14, p. e115. [\[CrossRef\]](#)

How to cite this article:

BI, Y.; ZHENG, X.; SHAO, C.; PAN, W.; JIANG, L. and OUYANG, H. (2011). Construction and application of a built-in dual luciferase reporter for microRNA functional analysis. *Electronic Journal of Biotechnology*, vol. 14, no. 2. <http://dx.doi.org/10.2225/vol14-issue2-fulltext-9>